



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 9/10, 15/54, 15/63, 15/70, 15/82, 5/10, A01H 5/00, C12P 21/02, 19/04, 19/18	A2	(11) International Publication Number: WO 00/47727 (43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: PCT/EP00/00954 (22) International Filing Date: 7 February 2000 (07.02.00) (30) Priority Data: 199 05 069.4 8 February 1999 (08.02.99) DE (71) Applicants (for all designated States except US): PLANT-TEC BIOTECHNOLOGIE GMBH FORSCHUNG & ENTWICKLUNG [DE/DE]; Hermannswerder 14, D-14473 Potsdam (DE). MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN e.V. [DE/DE]; Berlin (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): KOSSMANN, Jens [DE/DE]; Düppelstr. 29, D-14163 Berlin (DE). WELSH, Thomas [DE/DE]; Schönburgstrasse 20, D-12103 Berlin (DE). QUANZ, Martin [DE/DE]; Oppelner Strasse 34, D-10997 Berlin (DE). KNUTH, Karola [DE/DE]; Schöneberger Str. 6, D-12103 Berlin (DE). (74) Agent: VOSSIUS & PARTNER; Siebertstrasse 4, D-81765 München (DE).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NUCLEIC ACID MOLECULES ENCODING ALTERNANSUCRASE		
(57) Abstract <p>Nucleic acid molecules encoding an alternansucrase are provided. Moreover, vectors, host cells and plant cells transformed by the herein-described nucleic acid molecules and plants containing them are provided. Furthermore, methods are described for preparing transgenic plants which synthesize the carbohydrate alternan, because of the insertion of nucleic acid molecules encoding an alternansucrase. Moreover, methods for preparing alternan and products resulting from them are provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Nucleic Acid Molecules Encoding Alternansucrase

The present invention relates to nucleic acid molecules encoding an alternansucrase. Moreover, this invention relates to vectors, host cells and plant cells transformed with the herein-described nucleic acid molecules, and plants containing said cells. Moreover, methods for preparing transgenic plants which due to the insertion of DNA molecules encoding an alternansucrase, synthesize the carbohydrate alternan, are described. Furthermore, methods for preparing alternan are described.

Prior art documents, the disclosure content of which is included into the present application by reference thereto, are cited hereinafter.

Alternan is a polysaccharide composed of glucose units. The glucose units are linked to each other via α -1,3- and α -1,6-glycosidic bonds, and said two types of bonds predominantly appear alternatingly. However, alternan is not a linear polysaccharide, but may contain branches (Seymour et al., Carbohydrate Research 74, (1979), 41-62). Because of its physico-chemical properties, the possibilities of application of alternan both in the pharmaceutical industry, for instance as a carrier of pharmaceutically active ingredients and as an additive in the textile, cosmetics and food industry have been discussed (Lopez-Munguia et al., Enzyme Microb. Technol. 15, (1993), 77-85; Leathers et al., Journal of Industrial Microbiology & Biotechnology 18, (1997), 278-283). Moreover, it can be used as a substitute for gum arabic (Coté, Carbohydrate Polymers 19, (1992), 249-252).

Industry has a high interest in biotechnological methods for preparing oligosaccharides and polysaccharides, and in particular alternan which is hardly or not at all accessible to classical organic synthesis. Compared to the classical approach of organic synthesis chemistry, biotechnological processes offer

advantages. For instance, enzymatically catalyzed reactions as a rule show much higher specificities (regio specificity, stereo specificity) and higher reaction speeds, proceed under milder reaction conditions and lead to higher yields. These factors are of outstanding importance in the preparation of new oligosaccharides and polysaccharides.

Alternan is prepared enzymatically with the use of enzymes possessing the biological activity of alternansucrases. Alternansucrases belong to the group of glucosyltransferases, which, starting from saccharose, are able to catalyze the formation of alternan and fructose. So far, alternansucrases have only been found in the bacterium *Streptococcus mutans* (Mukasa et al. (J. Gen. Microbiol. 135 (1989), 2055-2063); Tsumori et al. (J. Gen. Microbiol. 131 (1985), 3347-3353)) and in specific strains of the gram positive bacterium *Leuconostoc mesenteroides* where they are, as a rule, present together with other polysaccharide-forming enzymes, such as for instance dextran-forming dextransucrases, or together with polysaccharide-degrading enzymes, such as alternanases. Hence, the naturally occurring strains also produce dextran in addition to alternan.

So far, alternan has been prepared in a cell-free system using partially purified proteins or by fermentation using alternansucrase-producing strains of *Leuconostoc mesenteroides*.

Various purification methods for the purification of alternansucrases have been described (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85; Lopez-Munguia et al., Annals New York Academy of Sciences 613 (1990), 717-722; Coté and Robyt, Carbohydrate Research 101 (1982), 57-74). These methods are complex and relatively costly, and, as a rule, lead to low protein yields (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283). None of these methods allows highly pure alternansucrase protein to be produced, and therefore sequencing of the protein and the isolation of the corresponding DNA sequences have not been successful so far. If the alternansucrase protein purified according to these methods is used for *in vitro* preparation of alternan, then the dextransucrase protein residues contained in the alternansucrase preparation produce dextran impurities in the alternan produced. The separation of alternan and dextran is relatively time-consuming and costly (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283). Another disadvantage of the

dextranase protein impurities contained in the enzyme preparation of alternanase protein is the fact that a part of the saccharose substrate is converted into dextran and not into alternan, which results in a reduction of the alternan yield.

The fermentative preparation by means of *Leuconostoc* also leads to the formation of product mixtures of alternan and dextran. In order to increase the amount of alternanase from *Leuconostoc* strains, mutants have been isolated, such as the mutant NRRL B-21138, which secrete the alternanase and lead to a higher proportion of the amount of alternanase formed relative to dextranase. However, if such mutants are fermented with sucrose, the alternan obtained continues to show dextran impurities (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283).

As can be seen from the prior art discussed above, it has not been possible to provide highly purified alternanase protein so far.

Hence, the present invention addresses the problem of providing means and methods allowing alternan to be prepared in a time-saving and inexpensive manner.

This problem is solved by the provision of the embodiments characterized in the patent claims.

Consequently, the present invention relates to a nucleic acid molecule encoding a protein possessing the biological activity of an alternanase selected from the group consisting of

- (a) nucleic acid molecules encoding at least the mature form of a protein which comprises the amino acid sequence indicated in Seq. ID No. 2 or the amino acid sequence encoded by the cDNA contained in plasmid DSM 12666;
- (b) nucleic acid molecules comprising the nucleotide sequence indicated in Seq. ID No. 1 or the nucleotide sequence of the cDNA contained in plasmid DSM 12666 or a corresponding ribonucleotide sequence;

- (c) nucleic acid molecules encoding a protein, the amino acid sequence of which has a homology of at least 40% to the amino acid sequence indicated in Seq. ID No. 2;
- (d) nucleic acid molecules, one strand of which hybridizes with the nucleic acid molecules as defined in (a) or (b);
- (e) nucleic acid molecules comprising a nucleotide sequence encoding a biologically active fragment of the protein which is encoded by any one of the nucleic acid molecules as defined in (a), (b), (c) or (d); and
- (f) nucleic acid molecules, the nucleotide sequence of which deviates because of the degeneration of the genetic code from the sequence of the nucleic acid molecules as defined in (a), (b), (c), (d) or (e).

Consequently, the present invention relates to nucleic acid molecules encoding proteins possessing the biological activity of an alternansucrase, said molecules preferably encoding proteins comprising the amino acid sequence indicated in Seq. ID No. 2.

An enzyme possessing the enzymatic or biological activity of an alternansucrase (E.C. 2.4.1.140) is understood to mean an enzyme which is able to catalyze the conversion of saccharose into alternan and fructose. This conversion may occur both in the presence and absence of external acceptors (for instance maltose, isomaltose, isomaltotriose etc.). In the absence of external acceptors, alternansucrases starting from saccharose catalyze the release of fructose and high molecular alternan, a polysaccharide composed of glucose units, the backbone of which consists of glucose units predominantly connected to each other alternatingly by α -1,3- and α -1,6-glycosidic bonds. Concerning the percentage of α -1,3- and α -1,6-linked glucose units the literature displays different values. According to Mukasa et al. (J. Gen. Microbiol. 135 (1989), 2055-2063), alternan consists of 76 mol% α -1,3-linked glucose and 24 mol% α -1,6-linked glucose. Tsumori et al. (J. Gen. Microbiol. 131 (1985), 3347-3353) describe alternan as a polyglucan containing 49.1 mol% α -1,6-linked glucose and 33.9 mol% α -1,3-linked glucose with 13.6 mol% terminal glucose and 3.3 mol% α -1,3,6-branched glucose. In the presence of external acceptors, such as maltose, isomaltose, isomaltotriose and

methyl- α -D-glucan, alternansucrase can catalyze the synthesis of α -D-glucan chains, in which the glucose residues are predominantly alternatingly connected by α -1,6- and α -1,3-glycosidic bonds, and the synthesis of fructose at these polysaccharide acceptors. Depending on the acceptor used, the products formed have different structures. The enzymatic activity of an alternansucrase can for instance be detected as described by Lopez-Munguia (Annals New York Academy of Sciences 613 (1990), 717-722) or as described in the examples of the present application.

The invention in particular relates to nucleic acid molecules containing the nucleotide sequence indicated under Seq. ID No. 1 or a part thereof, and preferably to molecules, which comprise the coding region indicated in Seq. ID No. 1 or corresponding ribonucleotide sequences.

Moreover, the present invention relates to nucleic acid molecules which encode an alternansucrase and the one strand of which hybridizes with one of the above-described molecules.

The present invention also relates to nucleic acid molecules which encode a protein, which has a homology, that is to say an identity of at least 40%, preferably at least 60%, preferably at least 70%, especially preferably at least 80% and in particular at least 90% to the entire amino acid sequence indicated in Seq. ID No. 2, the protein possessing the biological activity of an alternansucrase.

The present invention also relates to nucleic acid molecules, which encode an alternansucrase and the sequence of which deviates on account of the degeneration of the genetic code from the nucleotide sequences of the above-described nucleic acid molecules.

The invention also relates to nucleic acid molecules possessing a sequence which is complementary to the whole or a part of the above-mentioned sequences.

The nucleic acid sequence indicated in Seq. ID No. 1 for instance encodes an extracellular alternansucrase. Secretion is ensured by a signal sequence which comprises the first approximately 39 N-terminal amino acid groups of the Seq. ID No. 2. In certain circumstances it may be desirable for only the mature protein to be expressed without naturally occurring signal sequences and/or together with other

signal sequences. Hence, the above-described nucleic acid molecules encode at least the mature form of a protein possessing the biological activity of an alternansucrase.

Within the present invention the term "hybridization" means hybridization under conventional hybridization conditions, preferably under stringent conditions, as for instance described in Sambrook et al., *Molecular Cloning, A Laboratory Manual*, 2nd edition (1989) Cold Spring Harbour Laboratory Press, Cold Spring Harbour, NY. Within an especially preferred meaning the term "hybridization" means that hybridization occurs under the following conditions:

Hybridization buffer: 2 x SSC; 10 x Denhardt solution (Fikoll 400 + PEG + BSA; ratio 1:1:1); 0.1% SDS; 5 mM EDTA; 50 mM Na₂HPO₄; 250 µg/ml of herring sperm DNA; 50 µg/ml of tRNA; or 0.25 M of sodium phosphate buffer, pH 7.2; 1 mM EDTA 7% SDS

Hybridization temperature T = 60°C

Washing buffer: 2 x SSC; 0.1% SDS

Washing temperature T = 60°C.

Nucleic acid molecules which hybridize with the nucleic acid molecules of the invention can, in principle, encode alternansucrases from any organism expressing such proteins.

Nucleic acid molecules which hybridize with the molecules of the invention can for instance be isolated from genomic libraries of microorganisms. Alternatively, they can be prepared by genetic engineering or chemical synthesis.

Such nucleic acid molecules may be identified and isolated with the use of the molecules of the invention or parts of these molecules or reverse complements of these molecules, for instance by hybridization according to standard methods (see for instance Sambrook et al., 1989, *Molecular Cloning. A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

Nucleic acid molecules possessing the same or substantially the same nucleotide sequence as indicated in Seq. ID No. 1 or parts thereof can, for instance, be used

as hybridization probes. The fragments used as hybridization probes can also be synthetic fragments which are prepared by usual synthesis techniques, and the sequence of which substantially coincides with that of an inventive nucleic acid molecule.

The molecules hybridizing with the nucleic acid molecules of the invention also comprise fragments, derivatives and allelic variants of the above-described nucleic acid molecules encoding an alternansucrase of the invention. Herein, fragments are understood to mean parts of the nucleic acid molecules which are long enough to encode one of the described proteins, preferably showing the biological activity of an alternansucrase. In this connection, the term derivative means that the sequences of these molecules also differ from the sequences of the above-described nucleic acid molecules in one or more positions and show a high degree of homology to these sequences. In this context, homology means a sequence identity of at least 40%, in particular an identity of at least 60%, preferably more than 80% and particularly preferably more than 90%. Deviations from the above-described nucleic acid molecules may have been produced by deletion, substitution, insertion and/or recombination.

Preferably, the degree of homology is determined by comparing the respective sequence with the nucleotide sequence of the coding region of SEQ ID No.1. When the sequences which are compared do not have the same length, the degree of homology preferably refers to the percentage of nucleotide residues in the shorter sequence which are identical to nucleotide residues in the longer sequence. The degree of homology can be determined conventionally using known computer programs such as the ClustalW program (Thompson et al., Nucleic Acids Research 22 (1994), 4673-4680) distributed by Julie Thompson (Thompson@EMBL-Heidelberg.DE) and Toby Gibson (Gibson@EMBL-Heidelberg.DE) at the European Molecular Biology Laboratory, Meyerhofstrasse 1, D 69117 Heidelberg, Germany. ClustalW can also be downloaded from several websites including IGBMC (Institut de Génétique et de Biologie Moléculaire et Cellulaire, B.P.163, 67404 Illkirch Cedex, France; <ftp://ftp-igbmc.u-strasbg.fr/pub/>) and EBI (<ftp://ftp.ebi.ac.uk/pub/software/>) and all sites with mirrors to the EBI (European

Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SD, UK).

When using ClustalW program version 1.8 to determine whether a particular sequence is, for instance, 90% identical to a reference sequence according to the present invention, the settings are set in the following way for DNA sequence alignments:

KTUPLE=2, TOPDIAGS=4, PAIRGAP=5, DNAMATRIX:IUB, GAOPEN=10, GAPEXT=5, MAXDIV=40, TRANSITIONS: unweighted.

For protein sequence alignments using ClustalW program version 1.8 the settings are the following: KTUPLE=1, TOPDIAG=5, WINDOW=5, PAIRGAP=3, GAOPEN=10, GAPEXTEND=0.05, GAPDIST=8, MAXDIV=40, MATRIX=GONNET, ENDGAPS(OFF), NOPGAP, NOHGAP.

Furthermore, homology means preferably that the encoded protein displays a sequence identity of at least 40%, more preferably of at least 60%, even more preferably of at least 80%, in particular of at least 90% and particularly preferred of at least 95% to the amino acid sequence depicted under SEQ ID NO: 2.

Homology, moreover, means that there is a functional and/or structural equivalence between the corresponding nucleic acid molecules or proteins encoded thereby. Nucleic acid molecules which are homologous to the above-described molecules and represent derivatives of these molecules are, as a rule, variations of these molecules which represent modifications having the same biological function. They may be either naturally occurring variations, for instance sequences from other microorganisms, or mutations, and said mutations may have formed naturally or may have been produced by deliberate mutagenesis. Furthermore, the variations may be synthetically produced sequences. The allelic variants may be naturally occurring variants or synthetically produced variants or variants produced by recombinant DNA techniques.

In a further preferred embodiment the term "derivative" encompasses a nucleic acid molecule coding for a protein which comprises at least one, more preferably at least three, even more preferably at least five, in particular at least ten and particularly preferred at least twenty of the peptide motifs selected from the group consisting of

- a) MKQQE (SEQ ID NO: 22),
- b) KKVPV (SEQ ID NO: 23),

- c) KDDEN (SEQ ID NO: 24),
- d) IDGNL (SEQ ID NO: 25),
- e) YVADS (SEQ ID NO: 26),
- f) HLRKN (SEQ ID NO: 27),
- g) NENTP (SEQ ID NO: 28),
- h) NVDGY (SEQ ID NO: 29),
- i) NPDLK (SEQ ID NO: 30),
- j) SNDSG (SEQ ID NO: 31),
- k) NTFVK (SEQ ID NO: 32),
- l) ISGYL (SEQ ID NO: 33),
- m) SNAAL (SEQ ID NO: 34),
- n) RQYTD (SEQ ID NO: 35),
- o) QLYRA (SEQ ID NO: 36),
- p) DDKAP (SEQ ID NO: 37),
- q) TRQYT (SEQ ID NO: 38),
- r) ITFAG (SEQ ID NO: 39),
- s) NQYKG (SEQ ID NO: 40),
- t) LFLNA (SEQ ID NO: 41),
- u) QVSDT (SEQ ID NO: 42),
- v) LITLN (SEQ ID NO: 43),
- w) GRYVH (SEQ ID NO: 44),
- x) TAPYG (SEQ ID NO: 45),
- y) VVDYQ (SEQ ID NO: 46),
- z) LSGQE (SEQ ID NO: 47).

The proteins encoded by the different variants of the nucleic acid molecules of the invention possess certain characteristics they have in common. These include for instance enzymatic activity, molecular weight, immunological reactivity, conformation, etc., and physical properties, such as for instance the migration behavior in gel electrophoreses, chromatographic behavior, sedimentation coefficients, solubility, spectroscopic properties, stability, pH optimum, temperature optimum etc.

Alternansucrase (E.C. 2.4.1.140) is an enzyme belonging to the group of glucosyltransferases. So far, alternansucrase activity has not been found in plants, but only in the bacterium *Streptococcus mutans* (Mukasa et al. (J. Gen. Microbiol. 135 (1989), 2055-2063); Tsumori et al. (J. Gen. Microbiol. 131 (1985), 3347-3353)) and in specific strains of the bacterium *Leuconostoc mesenteroides*, for instance in NRRL B-1355, NRRL B-1498 and NRRL B-1501. As a rule, these strains contain different glucosyltransferases and secrete dextransucrases apart from alternansucrases if they are allowed to grow on saccharose-containing media. As a rule, these two sucrases possess a high binding affinity to the polysaccharides synthesized by them (Lopez-Munguia et al., Annals New York Academy of Sciences 613 (1990), 717-722) with the result that these polysaccharides must be separated from the protein in the purification of the enzymes from *Leuconostoc mesenteroides* strains grown on saccharose-containing medium (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85; Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283).

In the absence of external acceptors, alternansucrases, starting from saccharose, catalyze the release of fructose and high molecular alternan, a polysaccharide which is composed of glucose units, and the backbone of which consists of glucose units predominantly linked to each other alternately by α -1,3- and α -1,6-glycosidic bonds and which according to light scattering measurement data should have a molecular weight of $>10^7$ (Coté, Carbohydrate Polymer 19 (1992), 249-252). To date there has been no report of alternan possessing a terminal fructose residue. Nevertheless, the existence of a terminal fructose unit in alternan can not be completely excluded. Lopez-Munguia et al. (Enzyme Microb. Technol. 15 (1993) 77-85) describe that alternan is resistant to degradation by dextranases. However, it can be degraded by so-called alternanases, whereby ring-shaped oligomers of alternan of different polymerization degree can be produced (Biely et al., Eur. J. Biochem. 226 (1994), 633-639). Ultrasonic treatment of high molecular alternan allows the molecular weight of alternan to be reduced to $<10^6$ (Coté, Carbohydrate Polymers 19 (1992), 249-252). If aqueous solutions of this ultrasonically treated alternan are prepared, then these solutions show rheological properties comparable to those of aqueous solutions of gum arabic. So-called "limit alternan" having a molecular weight of about 3500 can be produced by enzymatic degradation using

isomaltodextranase from *Arthrobacter globiformis* (NRRL B-4425) (Coté, Carbohydrate Polymers 19 (1992), 249-252).

In the presence of external acceptors, such as for instance maltose, isomaltose, isomaltotriose and methyl- α -D-glucan, alternansucrase catalyzes at said saccharide acceptors the synthesis of α -D-glucan chains, in which the glucose moieties are predominantly alternatingly linked by α -1,6- and α -1,3 glycosidic bonds, and the synthesis of fructose. Depending on the acceptor used, the resulting products have different structures and a molecular weight which is lower than that of high molecular alternan and a polymerization degree of <15 . Because of the polymerization degree, these products are often also referred to as oligoalternans (Pelenc et al., Sciences Des Aliments 11 (1991), 465-476). However, within the framework of the present invention these low molecular products which can be prepared in the presence of external acceptors are also to be referred to as alternan.

In the preparation of oligoalternans by means of partially purified alternansucrase protein, maltose is an acceptor (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85) producing high oligoalternan yields. Panose (degree of polymerization (d.p.) of 3) is the first acceptor product which is formed starting from maltose through the formation of an α -1,6-glycosidic bond.

In contrast thereto, isomaltose is a less effective acceptor which leads to lower yields of oligoalternan (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85).

Alternansucrase is relatively stable and has a half life period of 2 days in 50 mM of acetate buffer, pH 5.4 at 40°C (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85). The enzyme shows maximum activity at a temperature of 40°C and a pH value of 5.6 (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85).

In the absence of the substrate saccharose, alternansucrase catalyzes disproportionation reactions leading to a (partial) rearrangement of alternan. In particular when partially purified alternansucrase preparations containing dextransucrase contaminations were used to prepare oligoalternans, high disproportionation rates were found which lead to a complete rearrangement of oligoalternan (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85).

For the molecular weight of alternansucrase according to SDS PAGE determination, different numerical values can be found: 135 kDa, 145 kDa, 173 kDa and 196 kDa, respectively (Leathers et al., Journal of Industrial Microbiology & Biotechnology 18 (1997), 278-283; Kim & Robyt, Enzyme Microb. Technol. 16 (1994), 659-664; Zhanley & Smith, Applied and Environmental Microbiology 61(3) (1995), 1120-1123).

The enzymatic activity of an alternansucrase can be shown for instance as described in Lopez-Munguia et al. (Annals New York Academy of Sciences 613 (1990), 717-722) or as described in the examples of the present application.

One activity unit (1u) can be defined as the amount of enzyme leading to the release of 1 μ mol of fructose within one minute.

The nucleic acid molecules of the invention can be DNA molecules, in particular genomic molecules. Moreover, the nucleic acid molecules of the invention may be RNA molecules. The nucleic acid molecules of the invention can be obtained for instance from natural sources or may be produced synthetically or by recombinant techniques.

The nucleic acid molecules of the invention allow host cells to be prepared which produce recombinant alternansucrase protein of high purity and/or in sufficient quantities, and genetically engineered plants possessing an activity of these enzymes leading to the formation of alternan *in planta*. Within the framework of the present invention the term "high purity" means that the protein according to the invention displays a degree of purity of at least 80%, preferably of at least 90%, even more preferably of at least 95%. Moreover, means and methods are provided which may be used for preparing alternan using host cells and/or for preparing recombinant alternansucrase protein. Consequently, the provision of the nucleic acid molecules of the invention permits the preparation of alternan of high purity by methods which are relatively inexpensive and consume relatively little time.

In a preferred embodiment, the nucleic molecules of the invention are derived from microorganisms, preferably from bacteria, more preferably from gram-positive bacteria and in particular preferably from bacteria belonging to the genus

Leuconostoc. Nucleic acid molecules from bacteria belonging to the species *Leuconostoc mesenteroides* are particularly preferred.

The invention also relates to oligonucleotides specifically hybridizing to a nucleic acid molecule of the invention. Such oligonucleotides have a length of preferably at least 10, in particular at least 15, and particularly preferably of at least 50 nucleotides. They are characterized in that they specifically hybridize to the nucleic acid molecules of the invention, that is to say that they do not or only to a very minor extent hybridize to nucleic acid sequences encoding other proteins, in particular other glucosyltransferases. The oligonucleotides of the invention can be used for instance as primers for amplification techniques such as the PCR reaction or as a hybridization probe to isolate related genes.

Moreover, the invention relates to vectors, in particular plasmids, cosmids, viruses, bacteriophages and other vectors commonly used in gene technology, which contain the above-described nucleic acid molecules of the invention. In a preferred embodiment of the invention, the vectors of the invention lend themselves to the transformation of fungal cells or cells of microorganisms. Preferably, such vectors are suitable to transform plant cells. Particularly preferably, such vectors permit the integration of the nucleic acid molecules of the invention, possibly together with flanking regulatory regions, into the genome of the plant cell. Examples thereof are binary vectors which can be used in the *Agrobacteria*-mediated gene transfer, and some are already commercially available.

In another preferred embodiment, the nucleic acid molecules contained in the vectors are connected to regulatory elements ensuring the transcription and synthesis of a translatable RNA in prokaryotic or eukaryotic cells.

The expression of the nucleic acid molecules of the invention in prokaryotic or eukaryotic cells, for instance in *Escherichia coli*, is interesting because it permits a more precise characterization of the enzymatic activities of the enzymes encoded by these molecules. Moreover, it is possible to express these enzymes in such prokaryotic or eukaryotic cells which are free from interfering enzymes, such as dextranases or other polysaccharide-forming or polysaccharide-degrading

enzymes. In addition, it is possible to insert different mutations into the nucleic acid molecules by methods usual in molecular biology (see for instance Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY), leading to the synthesis of proteins possibly having modified biological properties. On the one hand it is possible in this connection to produce deletion mutants in which nucleic acid molecules are produced by progressive deletions from the 5' or 3' end of the coding DNA sequence, and said nucleic acid molecules lead to the synthesis of correspondingly shortened proteins. Such deletions at the 5' end of the nucleotide sequence for instance allow amino acid sequences to be identified which are responsible for the secretion of the enzyme in microorganisms (transit peptides).

This permits the deliberate preparation of enzymes which are no longer secreted by the removal of the corresponding sequences, but remain within the cell of the corresponding host organism or are localized in other compartments, for instance in the plastids, mitochondria, vacuole, on account of the addition of other signal sequences.

On the other hand, the introduction of point mutations is also conceivable at positions at which a modification of the amino acid sequence for instance influences the enzyme activity or the control of the enzyme. In this manner, it is for instance possible to produce mutants which possess a modified stereo and regio selectivity or a modified K_m value or which are no longer subject to the control mechanisms normally existing in the cell and realized via an allosteric control or covalent modification.

Moreover, mutants possessing a modified substrate or product specificity can be prepared. Furthermore, it is possible to prepare mutants having a modified activity-temperature-profile.

Furthermore, in the case of expression in plants, the insertion of mutations into the nucleic acid molecules of the invention allows the gene expression rate and/or the activity of the proteins encoded by the nucleic acid molecules of the invention to be increased.

For genetic engineering in prokaryotic cells, the nucleic acid molecules of the invention or parts of these molecules can be introduced into plasmids which permit mutagenesis or sequence modification by recombination of DNA sequences.

Standard methods (see Sambrook et al., 1989, Molecular Cloning: A laboratory manual, 2nd edition, Cold Spring Harbor Laboratory Press, NY, USA) allow base exchanges to be performed or natural or synthetic sequences to be added. DNA fragments can be connected to each other by applying adapters and linkers to the fragments. Moreover, engineering measures which provide suitable restriction sites or remove surplus DNA or restriction sites can be used. In those cases, in which insertions, deletions or substitutions are possible, *in vitro* mutagenesis, "primer repair", restriction or ligation can be used. In general, a sequence analysis, restriction analysis and other methods of biochemistry and molecular biology are carried out as analysis methods.

Moreover, the invention relates to plasmid pAlsu-pSK (see Fig. 2 and Example 2) which was deposited at Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ), Braunschweig, under the accession No. DSM 12666 on February 4, 1999, and to the nucleic acid molecules contained in the insert of plasmid DSM 12666 and encoding a protein possessing the enzymatic activity of an alternansucrase. Moreover, the present invention also relates to nucleic acid molecules which hybridize to the insertion of plasmid DSM 12666. Also, the present invention relates to nucleic acid molecules the nucleotide sequence of which deviates from that of the nucleic acid molecules of the plasmid DSM 12666 insert, because of the degeneration of the genetic code. Furthermore, the present invention relates to nucleic acid molecules which have a homology, that is to say a sequence identity of at least 40%, preferably of at least 60%, more preferably of at least 80%, even more preferably of at least 90%, and most preferably of at least 95% to the sequence of the insertion of plasmid DSM 12666.

Another embodiment of the invention relates to host cells, in particular prokaryotic or eukaryotic cells transformed with an above-described nucleic acid molecule of the invention or with a vector of the invention, and to cells descended from such transformed cells and containing a nucleic acid molecule or vector of the invention. According to another preferred embodiment, the host cells are cells of microorganisms. In the context of the present invention, the term "microorganism" comprises bacteria and all protists (e.g. fungi, in particular yeasts, algae) as defined

Schlegel's "Allgemeine Mikrobiologie" (Georg Thieme Verlag, 1985, 1-2). A preferred embodiment of the invention relates to cells of algae and host cells belonging to the genera *Aspergillus*, *Bacillus*, *Saccharomyces* or *Pichia* (Rodriguez, Journal of Biotechnology 33 (1994), 135-146, Romanos, Vaccine, Vol. 9 (1991), 901 et seq.). A particularly preferred embodiment of the invention relates to *E. coli* cells. Alternansucrase is especially preferably secreted by the host cell. The preparation of such host cells for the production of recombinant alternansucrase can be carried out by methods known to a man skilled in the art.

In a preferred embodiment of the invention, the host cells of the invention show no interfering enzymatic activities, such as those of polysaccharide-forming and/or polysaccharide-degrading enzymes.

An overview of different expression systems is for instance contained in Methods in Enzymology 153 (1987), 385-516, in Bitter et al. (Methods in Enzymology 153 (1987), 516-544) and in Sawers et al. (Applied Microbiology and Biotechnology 46 (1996), 1-9), Billman-Jacobe (Current Opinion in Biotechnology 7 (1996), 500-4), Hockney (Trends in Biotechnology 12 (1994), 456-463), Griffiths et al, Methods in Molecular Biology 75 (1997), 427-440). An overview of yeast expression systems is for instance given by Hensing et al. (Antonie van Leeuwenhoek 67 (1995), 261-279), Bussineau et al. (Developments in Biological Standardization 83 (1994), 13-19), Gellissen et al. (Antonie van Leeuwenhoek 62 (1992), 79-93, Fleer (Current Opinion in Biotechnology 3 (1992), 486-496), Vedvick (Current Opinion in Biotechnology 2 (1991), 742-745) and Buckholz (Bio/Technology 9 (1991), 1067-1072).

Expression vectors have been widely described in the literature. As a rule, they contain not only a selection marker gene and a replication-origin ensuring replication in the host selected, but also a bacterial or viral promoter, and in most cases a termination signal for transcription. Between the promoter and the termination signal there is at least one restriction site or a polylinker which enables the insertion of a coding DNA sequence. The DNA sequence naturally controlling the transcription of the corresponding gene can be used as the promoter sequence, if it is active in the selected host organism. However, this sequence can also be exchanged for other promoter sequences. It is possible to use promoters producing a constitutive expression of the gene and inducible promoters which permit a deliberate control of the expression of the postconnected gene. Bacterial and viral

promoter sequences possessing these properties are described in detail in the literature. Regulatory sequences for the expression in microorganisms (for instance *E. coli*, *S. cerevisiae*) are sufficiently described in the literature. Promoters permitting a particularly high expression of the postconnected gene are for instance the T7 promoter (Studier et al., Methods in Enzymology 185 (1990), 60-89), lacUV5, trp, trp-lacUV5 (DeBoer et al., in Rodriguez and Chamberlin (Eds), Promoters, Structure and Function; Praeger, New York, (1982), 462-481; DeBoer et al., Proc. Natl. Acad. Sci. USA (1983), 21-25), lp1, rac (Boros et al., Gene 42 (1986), 97-100). As a rule, the protein amounts are highest from the middle up to about the end of the logarithmic phase of the growth cycle of the microorganisms. Therefore, inducible promoters are preferably used for the synthesis of proteins. These promoters often lead to higher protein yields than do constitutive promoters. The use of highly constitutive promoters leads to the continuous transcription and translation of a cloned gene and thus often has the result that energy is lost for other essential cells functions with the effect that cell growth is slowed down (Bernard R. Glick/Jack J. Pasternak, Molekulare Biotechnologie (1995). Spektrum Akademischer Verlag GmbH, Heidelberg, Berlin, Oxford, p. 342). Therefore, in order to obtain an optimum amount of protein, a two-stage process is often used. First, the host cells are cultured under optimum conditions up to a relatively high cell density. In the second step, transcription is then induced depending on the type of promoter used. In this connection, a tac promoter is particularly suitable which can be induced by lactose or IPTG (=isopropyl- β -D-thiogalactopyranoside) (deBoer et al., Proc. Natl. Acad. Sci. USA 80 (1983), 21-25). Termination signals for transcription are also described in the literature.

The transformation of the host cell with DNA encoding an alternansucrase can, as a rule, be carried out by standard methods, as for instance described in Sambrook et al., (Molecular Cloning: A Laboratory Course Manual, 2nd edition (1989) Cold Spring Harbor Press, New York; Methods in Yeast Genetics, A Laboratory Course Manual, Cold Spring Harbor Laboratory Press, 1990). The host cell is cultured in nutrient media meeting the requirements of the particular host cell used, in particular in respect of the pH value, temperature, salt concentration, aeration, antibiotics, vitamins, trace elements etc.

Moreover, the invention relates to proteins and biologically active fragments thereof, which are encoded by the nucleic acid molecules of the invention and to methods for their preparation, wherein a host cell according to the invention is cultured under conditions permitting the synthesis of the protein, and the protein is subsequently isolated from the cultured cells and/or the culture medium.

According to a preferred embodiment of the invention, the alternansucrase is a recombinantly produced protein. In the context of the present invention, this is a protein prepared by inserting a DNA sequence encoding the protein into a host cell and expressing it therein. The protein can then be isolated from the host cell and/or the culture medium.

The nucleic acid molecules of the invention now allow host cells to be prepared which produce recombinant alternansucrase protein of high purity and/or in sufficient amounts. Within the framework of the present invention the term "high purity" means that the protein according to the invention displays a degree of purity of at least 80%, preferably of at least 90%, even more preferably of at least 95%. The time-consuming and costly methods already mentioned above, whereby alternansucrase protein which to date can only be obtained from particular *Leuconostoc* strains can be purified from other components such as for instance dextransucrases, polysaccharides, are dispensed with, because alternansucrase can be produced in host cells not possessing any adverse polysaccharide-synthesizing activities. Moreover, host cells and vectors can also be used, which allow the alternansucrase protein to be produced in the absence of saccharose, with the result that an additional separation of the alternansucrase protein from polysaccharides is no longer necessary. Moreover, the selection of suitable host cells and vectors allows alternansucrase protein to be provided in sufficient amounts, which has not been possible with the systems so far described.

Alternansucrase produced by the host cells can be purified by conventional purification methods, such as precipitation, ion exchange chromatography, affinity-chromatography, gel filtration, HPLC Reverse Phase Chromatography etc.

The modification of the nucleic acid molecules of the invention encoding an alternansucrase and expressed in the host cells, allows to produce a polypeptide in the host cell which is easier to isolate from the culture medium because of particular

properties. Thus, the protein to be expressed can be expressed as a fusion protein with an additional polypeptide sequence, the specific binding properties of which permit the isolation of the fusion protein by affinity chromatography (e.g. Hopp et al., Bio/Technology 6 (1988), 1204-1210; Sassenfeld, Trends Biotechnol. 8 (1990), 88-93).

Another embodiment of the invention relates to proteins possessing the enzymatic activity of an alternansucrase, in particular that from microorganisms, preferably Gram-positive microorganisms, particularly microorganisms of the genus *Leuconostoc*, and particularly preferably that from *Leuconostoc mesenteroides*. The molecular weight of the protein indicated in Seq. ID No. 2, as determined by calculation, is 228.96 kDa. The invention also relates to alternansucrases which possess a molecular weight of $229 \text{ kDa} \pm 120 \text{ kDa}$, preferably $229 \text{ kDa} \pm 50 \text{ kDa}$, and particularly preferably $230 \text{ kDa} \pm 25 \text{ kDa}$. The molecular weight of the mature protein, as determined by calculation, is 224.77 kDa.

The provision of the nucleic acid molecules of the invention, for the first time, makes it possible to prepare alternansucrase-expressing plant cells by means of genetic engineering, which was not possible so far, because classical culturing methods do not allow bacterial and fungal genes to be expressed in plants.

The invention, therefore, also relates to transgenic plant cells transformed by a nucleic acid molecule of the invention or a vector of the invention or descended from such cells, the nucleic acid molecule which encodes the protein that has the biological activity of an alternansucrase being under the control of regulatory elements permitting the transcription of a translatable mRNA in plant cells.

The introduction of the activity of the proteins of the invention, for instance by expression of corresponding nucleic acid molecules, opens the possibility of producing alternan in plant cells correspondingly modified by genetic engineering. Hence, the expression of the nucleic acid molecules of the invention in plant cells is possible, allowing an additional, corresponding alternansucrase activity not present in the wild type to be introduced. Moreover, it is possible to modify the nucleic acid molecules of the invention according to methods known to a skilled person, in order to obtain alternansucrases of the invention which for instance possess modified

temperature dependencies or substrate or product specificities. Such methods have already been described in more detail in a different context above.

A plurality of techniques is available by which DNA can be inserted into a plant host cell. These techniques include the transformation of plant cells by T-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as a transforming agent, the fusion of protoplasts, injection, electroporation of DNA, insertion of DNA by the biolistic approach and other possibilities.

The use of the Agrobacteria-mediated transformation of plant cells has been extensively investigated and sufficiently described in EP 120 516; Hoekema, In: The Binary Plant Vector System, Offsetdrukkerij Kanters B.V., Alblasterdam (1985), Chapter V; Fraley et al, Crit. Rev. Plant Sci. 4 (1993), 1-46 and An et al., EMBO J. 4 (1985), 277-287. Regarding the transformation of potatoes see for instance Rocha-Sosa et al. (EMBO J. 8 (1989), 29-33).

The transformation of monocotyledonous plants by means of Agrobacterium-based vectors has also been described (Chan et al., Plant Mol. Biol. 22 (1993), 491-506; Hiei et al., Plant J. 6 (1994) 271-282; Deng et al, Science in China 33 (1990), 28-34; Wilmink et al, Plant Cell Reports 11 (1992), 76-80; May et al., Bio/Technology 13 (1995), 486-492; Conner and Dormisse, Int. J. Plant Sci. 153 (1992), 550-555; Ritchie et al. Transgenic Res. 2 (1993), 252-265). An alternative system for transforming monocotyledonous plants is the transformation by the biolistic approach (Wan and Lemaux, Plant Physiol. 104 (1994), 37-48; Vasil et al., Bio/Technology 11 (1993), 1553-1558; Ritala et al., Plant Mol. Biol. 24 (1994) 317-325; Spencer et al., Theor. Appl. Genet. 79 (1990), 625-631), protoplast transformation, electroporation of partially permeabilized cells, insertion of DNA via glass fibers. The transformation of maize in particular has been repeatedly described in the literature (see for instance WO 95/06128, EP 0 513 849, EP 0 465 875, EP 29 24 35; Fromm et al, Biotechnology 8, (1990), 833-844; Gordon-Kamm et al., Plant Cell 2, (1990), 603-618; Koziel et al., Biotechnology 11 (1993), 194-200; Moroc et al., Theor. Appl. Genet. 80, (1990), 721-726).

The successful transformation of other types of cereals has also been described for instance of barley (Wan and Lemaux, supra; Ritala et al., supra, Krens et al., Nature 296 (1982), 72-74) and wheat (Nehra et al., Plant J. 5 (1994), 285-297). Generally, any promoter active in plant cells is suitable to express the nucleic acid

molecules in plant cells. The promoter can be so chosen that the expression in the plants of the invention occurs constitutively or only in a particular tissue, at a particular time of plant development or at a time determined by external influences. The promoter may be homologous or heterologous to the plant.

Suitable promoters are for instance the promoter of 35S RNA of the Cauliflower Mosaic Virus (see for instance US-A-5,352,605) and the ubiquitin-promoter (see for instance US-A-5,614,399) which lend themselves to constitutive expression, the patatin gene promoter B33 (Rocha-Sosa et al., EMBO J. 8 (1989), 23-29) which lends itself to a tuber-specific expression in potatoes or a promoter ensuring expression in photosynthetically active tissues only, for instance the ST-LS1 promoter (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84 (1987), 7943-7947; Stockhaus et al., EMBO, J. 8 (1989) 2445-2451), the Ca/b-promoter (see for instance US-A-5,656,496, US-A-5,639,952, Bansal et al., Proc. Natl. Acad. Sci. USA 89 (1992), 3654-3658) and the Rubisco SSU promoter (see for instance US-A-5,034,322; US-A-4,962,028) or the glutelin promoter from wheat which lends itself to endosperm-specific expression (HMW promoter) (Anderson, Theoretical and Applied Genetics 96, (1998), 568-576, Thomas, Plant Cell 2 (12), (1990), 1171-1180), the glutelin promoter from rice (Takaiwa, Plant Mol. Biol. 30(6) (1996), 1207-1221, Yoshihara, FEBS Lett. 383 (1996), 213-218, Yoshihara, Plant and Cell Physiology 37 (1996), 107-111), the shrunken promoter from maize (Maas, EMBO J. 8 (11) (1990), 3447-3452, Werr, Mol. Gen. Genet. 202(3) (1986), 471-475, Werr, Mol. Gen. Genet. 212(2), (1988), 342-350), the USP promoter, the phaseolin promoter (Sengupta-Gopalan, Proc. Natl. Acad. Sci. USA 82 (1985), 3320-3324, Bustos, Plant Cell 1 (9) (1989), 839-853) or promoters of zein genes from maize (Pedersen et al., Cell 29 (1982), 1015-1026; Quatroccio et al., Plant Mol. Biol. 15 (1990), 81-93). However, promoters which are only activated at a point in time determined by external influences can also be used (see for instance WO 93/07279). In this connection, promoters of heat shock proteins which permit simple induction may be of particular interest. Moreover, seed-specific promoters such as the USP promoter from *Vicia faba* which ensures a seed-specific expression in *Vicia faba* and other plants may be used (Fiedler et al., Plant Mol. Biol. 22 (1993), 669-679; Bäumlein et al., Mol. Gen. Genet. 225 (1991), 459-467). Moreover, fruit-specific promoters, such as described in WO 91/01373 may be used too.

Moreover, a termination sequence may be present, which serves to terminate transcription correctly and to add a poly-A-tail to the transcript, which is believed to have a function in the stabilization of the transcripts. Such elements are described in the literature (see for instance Gielen et al., EMBO J. 8 (1989), 23-29) and can be replaced at will.

Such cells can be distinguished from naturally occurring plant cells *inter alia* by the fact that they contain a nucleic acid molecule of the invention which does not naturally occur in these cells. Moreover, such transgenic plant cells of the invention can be distinguished from naturally occurring plant cells in that they contain at least one copy of the nucleic acid molecule of the invention stably integrated in their genome.

Moreover, the plant cells of the invention can preferably be distinguished from naturally occurring plant cells by at least one of the following features: If the inserted nucleic acid molecule of the invention is heterologous to the plant cell, then the transgenic plant cells are found to have transcripts of the inserted nucleic acid molecules of the invention. The latter can be detected for instance by Northern blot analysis. The plants cells of the invention preferably contain a protein encoded by an inserted nucleic acid molecule of the invention. This can be shown for instance by immunological methods, in particular by Western blot analysis.

Transgenic plant cells can be regenerated to whole plants according to methods known to a person skilled in the art.

The present invention also relates to the plants obtainable by regeneration of the transgenic plant cells of the invention. Furthermore, it relates to plants containing the above-described transgenic plant cells.

In most plants, the photoassimilates in the form of sugars formed during photosynthesis within a plant, i.e. mainly in the form of saccharose, are transported to the corresponding target organs. As saccharose is the substrate of the polymerization reaction of alternansucrase, all plants, both monocotyledonous and dicotyledonous can, in principle, be modified by the nucleic acid molecule of the invention in respect of alternansucrase expression.

The expression in plants of the nucleic acid molecules of the invention encoding a protein having the enzymatic activity of an alternansucrase can, for instance, be used to achieve a modification of the viscosity of the extracts possibly obtained from the plants, said modification being achieved by the synthesis of alternan. In this connection, for instance tomatoes are of interest. The expression of an alternansucrase in a tomato fruit leads to the synthesis of alternan and results in a modification of the viscosity of extracts obtained from these fruits for instance for the production of tomato puree or tomato ketchup.

The expression of the nucleic acid molecules of the invention is in particular advantageous in those organs of the plant which show a higher saccharose content or store saccharose. Such organs are for instance the beet of sugar beet or the cane of sugar cane. As these plants normally do not store any appreciable amounts of starch, the alternans synthesized by the alternansucrase from these plants could be isolated in the pure form.

The site where the biosynthesis of the saccharose in the plant cell occurs is the cytosol. The storage site, however, is the vacuole. During its transport into the storage tissue of the sugar beet or the potato or during its transport into the endosperm of seeds, the saccharose must pass the apoplast. Hence, all three compartments, i.e. the cytosol, the vacuole, the apoplast, lend themselves to the expression of the nucleic acid molecules for the synthesis of alternan. In addition, the plastids also lend themselves thereto, as could for instance be shown by the expression of bacterial fructosyl transferases in amyloplasts. Said fructosyl transferases which likewise require saccharose as a substrate, were able to mediate the formation of "amylofructan" in amyloplasts (Smeekens, Trends in Plant Science, Vol. 2, No. 8 (1997), 286-288).

In the case of starch-producing plants, such as potatoes and maize, where the starch biosynthesis and starch storage normally take place in the amyloplasts, an expression of the alternansucrase in apoplasts, in the cytosol or in the vacuole would lead to an additional synthesis of oligosaccharides and/or polysaccharides in these compartments, which can mean an overall increase in the yield.

As in the case of potatoes the starch synthesized in the amyloplasts can be separated from the alternan synthesized in the apoplast, in the cytosol or in the vacuole, the very same plant can be used to recover starch and alternan.

Moreover, transgenic potato and maize plants are known, the starch synthesis of which in the tubers and grains, respectively, is completely inhibited due to the inhibition of ADP-glucose-pyrophosphorylase by an antisense construct. In the case of potatoes, soluble sugars, in particular saccharose and glucose, accumulate instead, for instance in the tubers (Müller-Röber et al., EMBO J. 11 (1992), 1229-1238). Alternan can be prepared in the cytosol, the vacuole or apoplast of these plants by the expression of an alternansucrase which uses saccharose as a substrate.

Therefore in another embodiment of the invention the plant cells of the invention are further characterized by a reduced ADP glucose pyrophosphorylase (AGPase) activity compared to corresponding cells from wild-type plants.

DNA molecules encoding AGPase are well known to the person skilled in the art and described for example in Müller-Röber et al. (Mol. Gen. Genet. 224 (1) (1990), 136-146). By using DNA molecules encoding an AGPase it is possible to produce plants by means of recombinant DNA techniques (for example by an antisense, a ribozyme or a cosuppression approach) showing a reduced AGPase activity. Furthermore AGPase mutants, for example from maize (brittle-2 and shrunken-2), with reduced AGPase activity are known to the person skilled in the art.

The term "reduced" means preferably a reduction of AGPase activity of at least 10%, more preferably of at least 50% and even more preferably of at least 80% in comparison to corresponding wild-type cells.

The activity of an AGPase can be determined according to Müller-Röber et al. (Mol. Gen. Genet. 224 (1) (1990), 136-146) or to methods known to a person skilled in the art.

The reaction which is catalyzed by alternansucrase is distinguished by the fact that a glucose moiety is transferred directly from saccharose to an existing carbohydrate acceptor. By contrast, in the case of plants, the biosynthesis of linear glucans from saccharose, proceeds in such a way that the saccharose is first separated into glucose and fructose, which are then each converted into activated intermediate ADP-glucose. The glucose moiety is transferred by the enzyme starch-synthase from the ADP glucose to an already existing glucan, whereby ADP is released. The

conversion of saccharose into two ADP glucose molecules requires several energy consuming reactions. Therefore, the energy consumption of the reaction catalyzed by alternansucrase is substantially lower than the energy consumption in the synthesis of polysaccharides from saccharose in plant cells, which can lead to an increased yield of synthesized oligo and/or polysaccharides in plants containing the nucleic acid molecules of the invention.

In the expression of the nucleic acid molecules in plants there exists in principle the possibility that the synthesized protein can be localized in any compartment of the plant cell (e.g. in the cytosol, plastids, vacuole, mitochondria) or the plant (e.g. in the apoplast). In order to achieve the localization in a particular compartment, the coding region must, where necessary, be linked to DNA sequences ensuring localization in the corresponding compartment. The signal sequences used must each be arranged in the same reading frame as the DNA sequence encoding the enzyme.

In order to ensure the location in the plastids it is conceivable to use one of the following transit peptides: of the plastidic Ferredoxin: NADP⁺ oxidoreductase (FNR) of spinach which is enclosed in Jansen et al. (Current Genetics 13 (1988), 517-522). In particular, the sequence ranging from the nucleotides -171 to 165 of the cDNA Sequence disclosed therein can be used, which comprises the 5' non-translated region as well as the sequence encoding the transit peptide. Another example is the transit peptide of the waxy protein of maize including the first 34 amino acid residues of the mature waxy protein (Klösken et al., Mol. Gen. Genet. 217 (1989), 155-161). It is also possible to use this transit peptide without the first 34 amino acids of the mature protein. Furthermore, the signal peptides of the ribulose biphosphate carboxylase small subunit (Wolter et al., Proc. Natl. Acad. Sci. USA 85 (1988), 846-850; Nawrath et al., Proc. Natl. Acad. Sci. USA 91 (1994), 12760-12764), of the NADP malat dehydrogenase (Gallardo et al., Planta 197 (1995), 324-332), of the glutathione reductase (Creissen et al., Plant J. 8 (1995), 167-175) or of the R1 protein Lorberth et al. (Nature Biotechnology 16, (1998), 473-477) can be used.

In order to ensure the location in the vacuole it is conceivable to use one of the following transit peptides: the N-terminal sequence (146 amino acids) of the patatin

protein (Sonnewald et al., Plant J. 1 (1991), 95-106) or the signal sequences described by Matsuoka und Neuhaus, Journal of Experimental Botany 50 (1999), 165-174; Chrispeels und Raikhel, Cell 68 (1992), 613-616; Matsuoka und Nakamura, Proc. Natl. Acad. Sci. USA 88 (1991), 834-838; Bednarek und Raikhel, Plant Cell 3 (1991), 1195-1206; Nakamura und Matsuoka, Plant Phys. 101 (1993), 1-5 .

In order to ensure the location in the mitochondria it is for example conceivable to use the transit peptide described by Braun et al.(EMBO J. 11, (1992), 3219-3227).

In order to ensure the location in the apoplast it is conceivable to use one of the following transit peptides: signal sequence of the proteinase inhibitor II-gene (Keil et al., Nucleic Acid Res. 14 (1986), 5641-5650; von Schaewen et al., EMBO J. 9 (1990), 30-33), of the levansucrase gene from *Erwinia amylovora* (Geier and Geider, Phys. Mol. Plant Pathol. 42 (1993), 387-404), of a fragment of the patatin gene B33 from *Solanum tuberosum*, which encodes the first 33 amino acids (Rosahl et al., Mol. Gen. Genet. 203 (1986), 214-220) or of the one described by Oshima et al. (Nucleic Acid Res. 18 (1990), 181).

The nucleic acid sequence indicated in Seq. ID No. 1 encodes an extracellular alternansucrase. Secretion is ensured by a signal sequence comprising the first approximately 39 N-terminal amino acid residues of the Seq. ID No. 2.

The transgenic plants may, in principle, be plants of any plant species, that is to say they may be monocotyledonous and dicotyledonous plants. Preferably, the plants are useful plants cultivated by man for nutrition or for technical, in particular industrial, purposes. They are preferably starch-storing plants, for instance cereal species (rye, barley, oat, wheat, millet, sago etc.), rice, pea, marrow pea, cassava and potato; tomato, rape, soybean, hemp, flax, sunflower, cow pea or arrowroot, fiber-forming plants (e.g. flax, hemp, cotton), oil-storing plants (e.g. rape, sunflower, soybean) and protein-storing plants (e.g. legumes, cereals, soybeans). The invention also relates to fruit trees and palms. Moreover, the invention relates to forage plants (e.g. forage and pasture grasses, such as alfalfa, clover, ryegrass) and vegetable plants (e.g. tomato, lettuce, chicory) and ornamental plants (e.g. tulips, hyacinths). Sugar-storing and/or starch-storing plants are preferred. Sugar

cane and sugar beet, and potato plants, maize, rice, wheat and tomato plants are particularly preferred.

A further subject of the invention is a method for the production of transgenic plant cells and transgenic plants which in comparison to non-transformed wildtype cells / non-transformed wildtype plants synthesize alternan. In this method the expression and/or the activity of proteins encoded by the nucleic acid molecules of the invention is increased in comparison to corresponding wild-type cells / wildtype plants which do not show any alternansucrase expression and/or activity. In particular, such a method comprises the expression of a nucleic acid molecule according to the invention in plant cells. The nucleic acid molecule according to the invention is preferably linked to a promoter ensuring expression in plant cells. In a particularly preferred embodiment the method comprises the introduction of a nucleic acid molecule according to the invention into a plant cell and regeneration of a plant from this cell.

Such an increase in expression may, e.g., be detected by Northern blot analysis. The increase in activity may be detected by testing protein extracts for their alternansucrase activity derived from plant cells. The enzymatic activity of an alternansucrase can be measured, for instance, as described in Lopez-Munguia et al. (Annals New York Academy of Sciences 613, (1990), 717-722) or as described in the examples of the present application.

The invention also relates to propagation material of the plants of the invention. The term "propagation material" comprises those components of the plant which are suitable to produce offspring vegetatively or generatively. Suitable means for vegetative propagation are for instance cuttings, callus cultures, rhizomes or tubers. Other propagation material includes for instance fruits, seeds, seedlings, protoplasts, cell cultures etc. The preferred propagation materials are tubers and seeds. The invention also relates to harvestable parts of the plants of the invention such as, for instance, fruits, seeds, tubers or rootstocks.

Another embodiment of the invention relates to methods for preparing alternan which comprise the step of extracting and isolating alternan from a plant of the invention.

The extraction and isolation of alternan from a plant of the invention may be carried out by standard methods, such as precipitation, extraction and chromatographic methods.

Moreover, the present invention relates to alternan obtainable from a plant of the invention or from propagation material of the invention.

Moreover, the present invention relates to a method for preparing alternan and/or fructose, wherein a host cell of the invention secretes an alternansucrase into a saccharose-containing culture medium and alternan and/or fructose is/are isolated from the culture medium.

A preferred embodiment of the method of the invention uses an alternansucrase recombinantly produced and secreted by the host cell into the culture medium, thus avoiding the necessity of breaking up the cells and purifying the protein further, because the secreted protein can be obtained from the supernatant. The residual components of the culture medium can be removed by methods usual in processing technology, such as dialysis, reverse osmosis, chromatographic methods, etc. The same applies to the concentration of the protein secreted into the culture medium. The secretion of proteins by microorganisms is normally mediated by N-terminal signal peptides (signal sequence, leader-peptide, transit peptide). Proteins possessing this signal sequence are able to penetrate the cell membrane of the microorganism. A secretion of proteins can be achieved by adding the DNA sequence encoding this signal peptide to the corresponding region encoding the alternansucrase.

The natural signal peptide of the expressed alternansucrase is preferred, that of the alternansucrase from *Leuconostoc mesenteroides* NRRL B 1355 (see the first approximately 25 to 45 N-terminal amino acid residues of Seq. ID No. 2) is particularly preferred.

The signal peptide of α -CGTase from *Klebsiella oxytoca* M5A1 (Fiedler et al., J. Mol. Biol. 256 (1996), 279-291) or a signal peptide as encoded by the nucleotides 11529-11618 of the sequence available under the GenBank accession number X86014 is most preferred.

The preparation of alternan and/or fructose requires neither activated glucose derivatives nor co-factors, as are necessary in most synthesis reactions for polysaccharides occurring within the cells. Hence, alternansucrase-secreting microorganisms can be cultured in saccharose-containing medium, the secreted alternansucrase leading to a synthesis of alternan and fructose in the culture medium.

Contrary to host cells from *Leuconostoc mesenteroides*, which secrete alternansucrase by nature, the host cells used according to the invention have the advantage that they do not secrete proteins possessing adverse polysaccharide-synthesizing side reactions, such as dextransucrase, with the result that outside the host cell, apart from alternan, no other polysaccharides can be formed which, as a rule, can be separated from alternan only by costly and time-consuming procedures. Moreover, the host cells according to a preferred embodiment of the invention do not have any adverse polysaccharide-degrading side activities, which would otherwise lead to losses in the yield of the alternan produced.

The method of the invention yields fructose apart from alternan. Fructose can be used for the inexpensive isolation of so-called "high-fructose-containing syrups" (HFCS). Conventional methods for preparing fructose on the one hand provide for the enzymatic break down of saccharose by means of an invertase or for the break down of starch into glucose units, mostly brought about by acid hydrolysis, and for subsequent enzymatic conversion of the glucose into fructose by glucose isomerases. However, both methods lead to mixtures of glucose and fructose. The two components must subsequently be separated from each other by chromatographic methods.

The separation of the two reaction products of the method of the invention, or the separation of the reaction products from the substrate saccharose can be achieved for example with the use of membranes permitting the penetration of fructose, but not the penetration of saccharose and/or alternans. If continuous removal of

fructose via such a membrane is provided for, a more or less complete conversion of saccharose occurs.

The isolation of alternan and fructose can be carried out by standard methods or can be carried out as for instance described in the working examples.

According to one embodiment of the method, the host cells originate from microorganisms, preferably from *Escherichia coli*.

In another embodiment, the method of the invention works with fungal host cells, in particular cells of yeasts, such as *Saccharomyces cerevisiae*. Yeast cells producing alternan in saccharose-containing medium because of the enzymatic activity of an alternansucrase, cannot be readily used, as yeasts secrete an invertase which breaks down the extracellular saccharose. The yeasts take up the resulting hexoses via a hexose transporter. However, one yeast strain has been described (Riesmeier et al. EMBO J. 11 (1992), 4705-4713) which carries a defective *suc2* gene, and therefore cannot secrete invertase. Moreover, these yeast cells do not contain a transportation system able to import saccharose into the cells. If such a strain is so modified by means of the nucleic acid molecules of the invention that it secretes an alternansucrase into the culture medium, then fructose and alternan will be synthesized in saccharose-containing medium. The resulting fructose can subsequently be taken up by the yeast cells.

In another preferred embodiment of this method the host cell of the invention is present in an immobilized form.

As a rule, host cells are immobilized by inclusion of the cells in a suitable material, such as alginate, polyacrylamide, gelatin, cellulose or chitosan. However, adsorption or covalent binding of the cells to a carrier material is also possible (Brodellius and Mosbach, Methods in Enzymology Vol. 135 (1987), 222-230) An advantage of the immobilization of cells is that it allows substantially higher cell densities to be achieved than does culturing in liquid culture. This results in a higher productivity. Moreover, the costs for agitation and aeration of the culture decrease as do the costs for measures to maintain sterility. Another important aspect is the

possibility of a continuous alternan production with the result that unproductive phases regularly occurring in fermentation processes can be avoided or at least greatly reduced.

Another embodiment of the invention relates to a method for preparing alternan and/or fructose, wherein

- a) a saccharose-containing solution is contacted with a protein of the invention under conditions permitting the conversion of saccharose into alternan and/or fructose; and
- b) alternan and/or fructose is/are isolated from the solution.

In this embodiment, the invention thus relates to a method for preparing alternan and/or fructose *in vitro* by means of a cell-free enzyme preparation. In this case, microorganisms which for instance secrete alternansucrase are cultured up to the stationary phase in a saccharose-free medium permitting the formation of alternansucrase protein. After removing the cells from the culture medium by centrifugation, the secreted enzyme can be recovered from the supernatant. The enzyme can subsequently be added to saccharose-containing solutions in order to synthesize alternan and/or fructose. Compared to the above-described synthesis of alternan in a system not freed from cells, this method offers the advantage that the reaction conditions can be controlled better and the reaction products are substantially purer and easier to purify. The purification of the protein can be carried out as already described above.

A preferred embodiment of the method of the invention uses a purified alternansucrase. Purified alternansucrase is understood to mean an enzyme which is largely free from cell components of the cells in which the protein is synthesized and shows no contamination with proteins possessing polysaccharide-synthesizing activities (e.g. dextranases) or degrading activities, and/or no contamination with (polysaccharide) acceptors. The term "purified alternansucrase" preferably means an alternansucrase possessing a degree of purity of at least 70%, preferably at least 85%, and particularly preferably at least 95%.

The use of a purified protein for preparing alternan and/or fructose offers various advantages. Compared to methods working with partially purified protein extracts,

the reaction medium of the method of the invention does not contain any residues of the production strain (microorganism) which is used for the purification of the protein or for its preparation by genetic engineering.

Moreover, the use of the purified protein is advantageous for food and pharmaceutical industry applications. Thanks to the fact that the reaction medium is defined in its composition and freed from all unnecessary components, the product is likewise more precisely defined in respect of its components. In consequence of this, the procedure for obtaining food and pharmaceutical industry approval of these products produced by genetic engineering requires substantially less documentation, especially since these products should not show any traces of a transgenic microorganism.

Moreover, contrary to the so far described *in vitro* methods in cell free systems using partially purified alternansucrase preparations, the method of the invention using a purified alternansucrase has the advantage that it allows highly pure alternan to be prepared without the occurrence of dextransucrase and dextran contaminations, because of the high purity of the protein of the invention. Moreover, the method of the invention permits the production of alternan in high yields, without losses caused for instance by adverse side reactions of a dextransucrase, which would convert part of the substrate saccharose into undesired dextran, the separation of which from alternan would only be possible using time-consuming and expensive methods.

The method of the invention produces fructose in addition to alternan. The fructose can be used for the inexpensive recovery of so-called "high-fructose-containing syrups" (HFCS). The method of the invention yields products of high purity, because of the use of a purified alternansucrase. Hence, compared to conventional methods for preparing HFCS from maize starch, which comprise costly process steps for removing the buffer salts by ion exchange, (Crabb and Mitchinson, TIBTECH 15 (1997), 349-352) the method of the invention does not require an expensive purification of the fructose.

Another preferred embodiment of the method of the invention uses a recombinantly prepared alternansucrase.

According to another preferred embodiment, the enzyme possessing the enzymatic activity of an alternansucrase is immobilized on a carrier material.

Immobilization of the alternansucrase offers the advantage that the enzyme being the catalyst of the synthesis reaction can be easily recovered from the reaction mixture and reused several times. As the purification of enzymes is normally costly and time consuming, immobilization and reutilization of the enzymes allow for a substantial cost saving. Another advantage is the degree of purity of the reaction products not containing any residual protein.

There are many carrier materials available for the immobilization of proteins, and coupling to the carrier material can be made via covalent or non-covalent bonds (for an overview see: *Methods in Enzymology* 135, 136, 137). Widely used carrier materials include for instance agarose, alginate, cellulose, polyacrylamide, silica or nylon.

According to another embodiment of the invention, the alternansucrase (immobilized on a carrier material) is present between two membranes, one of which allows fructose, but not saccharose and alternan to penetrate, the other one of which allows saccharose, but not alternan to penetrate. The supply with substrate occurs through the membrane which allows saccharose to penetrate it. The synthesized alternan remains in the space between the two membranes and the released fructose can be continuously removed from the reaction equilibrium via the membrane which only allows fructose to penetrate it. Such an arrangement permits an efficient separation of the reaction products, and thus the production of pure fructose.

Moreover, the separation of fructose by ion exchange chromatography has been described ("Starch Hydrolysis Products, Worldwide Technology, Production, and Application", Edited by F. W. Schenck, R. E. Hebeda, (1992), VCH Publishers, Inc., New York).

Thus, the use of alternansucrases for preparing pure fructose on the one hand involves the advantage that the relatively inexpensive substrate saccharose can be used as the starting material, and on the other hand the fructose can be isolated easily from the reaction mixture without additional enzymatic conversions or chromatographic methods.

Moreover, the invention relates to methods for preparing alternan and/or fructose, wherein

- a) a saccharose-containing solution is contacted with a protein of the invention and acceptor molecules under conditions permitting the conversion of saccharose to alternan and/or fructose; and
- b) alternan and/or fructose is/are isolated from the solution.

Within the framework of the present invention an acceptor molecule is understood to mean a molecule at which an alternansucrase is able to catalyze a chain-extending reaction. The acceptor which can be added to the reaction mixture at the beginning of the reaction is preferably a carbohydrate or a carbohydrate derivative. The use of external acceptors leads to the production of low molecular products which are to be designated alternan in the context of the present invention. The carbohydrate acceptor is preferably an oligo or polysaccharide, in particular a branched polysaccharide, such as dextrin, glycogen or amylopectin, preferably a linear polysaccharide, and particularly preferably a saccharide selected from the group consisting of maltose, isomaltose, isomaltotriose and methyl- α -D-glucan. If an extension of the alternan chain at these acceptors occurs, then products are formed which have a higher molecular weight than the educt. Where maltose, isomaltose, isomaltotriose and methyl- α -D-glucan are used, one obtains products which have a lower molecular weight than the alternan that can be prepared in the absence of external carbohydrate acceptors.

The size of the molecular weight of the oligoalternans prepared depends on the saccharose/acceptor ratio used. For instance the degree of polymerization of the products increases as the saccharose/isomaltose ratio increases.

Moreover, the saccharose/acceptor ratio has an influence on the oligoalternan yield. For instance, the oligoalternan yield increases as the saccharose/isomaltose ratio decreases.

The hitherto described methods for producing oligoalternan with the use of alternansucrases which the authors claim have been purified (Pelenc et al., Sciences Des Aliments 11 (1991), 465-476) only yielded product mixtures of oligoalternan and oligodextran, in the presence of the carbohydrate acceptor maltose. In this case, the synthesis of oligodextran is presumably attributable to dextransucrase-contaminations of the alternansucrase preparation. Compared to

this method, the method of the invention offers the advantage that the use of recombinantly produced alternansucrase protein not containing any dextransucrase contaminants permits the preparation of oligoalternan without the simultaneous formation of oligodextran. Thus, the method of the invention makes it possible to provide oligoalternan, without requiring additional costly purification steps for separating oligodextran.

According to another preferred embodiment, the enzyme possessing the enzymatic activity of an alternansucrase is immobilized on a carrier material.

According to another preferred embodiment of the method of the invention, a recombinantly produced alternansucrase is used.

Moreover, the present invention relates to end products containing alternan. In this context, end products are understood to mean cosmetic products, preferably food products, fodder and particularly preferably pharmaceutical products.

Finally, the present invention relates to a method for preparing the afore-mentioned products comprising one of the above-described alternan manufacturing methods of the invention and the formulation of the thus obtained alternan in a form which is suitable for one of the afore-mentioned applications of the corresponding product.

These and other embodiments are disclosed and obvious to a skilled person and embraced by the description and the examples of the present invention. Additional literature regarding one of the above-mentioned methods, means and applications, which can be used within the meaning of the present invention, can be obtained from the state of the art, for instance from public libraries for instance by the use of electronic means. This purpose can be served inter alia by public databases, such as the "medline", which are accessible via internet, for instance under the address <http://www.ncbi.nlm.nih.gov/PubMed/medline.html>. Other databases and addresses are known to a skilled person and can be obtained from the internet, for instance under the address <http://www.lycos.com>. An overview of sources and information

regarding patents and patent applications in biotechnology is contained in Berks, TIBTECH 12 (1994), 352-364.

Description of the Figures:

Fig. 1:

Linear map of the entire sequence region which was cloned after the screening of a genomic library of *Leuconostoc mesenteroides* NRRL B 1355 by the corresponding overlapping fragments of the clones AS-19B1, AS-19B2, AS-28B and AS-29Ba.

Fig. 2:

Plasmid map pAlsu-pSK

Fig. 3:

HPLC chromatogram: Preparation of oligoalternan in the presence of maltose (Example 2).

Fig. 4:

Plasmid map pAlsu-pET24a

Fig. 5:

SDS PAGE with subsequent assay of sucrase activity (see Example 6)

The following protein extracts are used

- 1 + 2) *E. coli* BL21(DE3) containing pAlsu-pET24a-3
- 3 + 4) *E. coli* BL21(DE3) containing pAlsu-pET24a-7
- 5 + 6) *E. coli* BL21 (DE3) containing pAlsu-pET24a-21
- 7 + 8) *E. coli* BL21 (DE3) containing pET24a
- 1, 3, 5, 7) culture before induction with IPTG
- 2, 4, 6, 8) culture at the end of culturing

Fig. 6:

HPLC chromatogram of dextran T10

Fig. 7:

HPLC chromatogram of dextran T10 after dextranase digestion

Fig. 8:

HPLC chromatogram of oligoalternan

Fig. 9

HPLC chromatogram of oligoalternan after dextranase digestion.

Fig. 10

Map of the expression cassette including the polylinker of the plasmid pBinAR-N.

Fig. 11

Plasmid map pat-Alsu-Hyg.

Fig. 12

Plasmid map fnr-Alsu-Hyg.

Examples

Vectors used in the examples:

1. pBinAR-N

By the use of standard methods (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)) we introduced a different polylinker (see figure 10) between the 35S Promoter and the OCS-Terminator into the plasmid pBinAR (Höfgen und Willmitzer, Plant Science 66 (1990), 221-230). The resulting plasmid was called pBinAR-N.

2. pBinAR-Hyg-N

Via standard methods (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)) we isolated an EcoRI/HinDIII-fragment from pBinAR-N containing the 35S Promoter, the polylinker and the OCS-Terminator. This fragment was then ligated into the same restriction

sites of plasmid pBIB-Hyg (Becker, Nucleic Acids Research 18 (1990), 203). The resulting plasmid was called pBinAR-Hyg-N.

3. pBinAR-pat-Hyg

By using the oligonucleotides Sp-pat-5' and Sp-pat-3' (s. SEQ ID Nos. 48 and SEQ ID No. 49) we amplified DNA molecules coding for the leader peptide of the patatin protein from potato (see SEQ ID No. 50, which differs from the sequence used by Sonnewald et al. Plant J. 1 (1991), 95-106) via a PCR approach using plasmid pgT5 (Rosahl et al., Mol. Gen. Genet. 203 (1986), 214-220; Sonnewald et al., Plant J. 1 (1991), 95-106) as a template. The resulting PCR products were cut by the restriction enzymes XbaI and Sall and then ligated into the plasmid pBinAR-Hyg-N which was linearized before by using the restriction enzymes SpeI and Sall. The resulting plasmid was called pBinAR-pat-Hyg.

PCR conditions:

Buffer and polymerase from Boehringer Mannheim (Pwo Polymerase No. 1644947)

DNA	0,2 ng
10x Buffer + MgSO ₄	5 µl
dNTPs (je 10 mM)	1 µl
Primer Sp-pat-5'	120 nM
Primer Sp-pat-3'	120 nM
Pwo Polymerase	1,0 units
distilled water	ad 50 µl

Reaction conditions:

Step 1	95°C	2:30 min
Step 2	95°C	0:30 min
Step 3	64°C	0:30 min
Step 4	72°C	0:30 min
	(plus 1 sec per cycle)	
Step 5	72°C	5:00 min.

The steps 2 to 4 were repeated 35 times in a cyclical manner.

4. pBinAR-FNR-Hyg

By using the oligonucleotides Sp-fnr-5' and Sp-fnr-3 (see SEQ ID No. 51 and 52) we amplified DNA molecules coding for the transit peptide of the FNR protein from spinach via a PCR approach using plasmid p6SocFNR-15 (Jansen et al., Current Genetics 13, (1988), 517-522) as a template. The resulting PCR products were cut by XbaI and Sall and then cloned into the SpeI/Sall-opened pBinAR-Hyg-N. Resulting plasmid was called pBinAR-fnr-Hyg.

PCR conditions:

Buffer and polymerase from Gibco BRL (Platinum Taq DNA Polymerase High Fidelity No. 1304-011)

DNA	0,2 ng
10x Buffer	5 µl
MgSO ₄	2,0 µl
dNTPs (per 10mM)	1 µl
Primer Sp-fnr-5'	150 nM
Primer Sp-fnr-3'	150 nM
Taq Platinum Hifi Polymerase	1,5 units
distilled water	ad 50 µl

Reaction conditions:

Step 1	95°C	2:30 min
Step 2	95°C	0:30 min
Step 3	58°C	0:30 min
Step 4	68°C	0:20 min
	(plus 1 sec per cycle)	
Step 5	68°C	3:00 min

The steps 2 to 4 were repeated 35 times in a cyclical manner.

Example 1: Cloning of alternansucrase from *Leuconostoc mesenteroides* NRRL-B1355Isolation and sequencing of alternansucrase

The strain *Leuconostoc mesenteroides* NRRL-B1355 was cultured in 1 l of Lactobacilli MRS Broth (Difco) complemented with 5% saccharose at 28°C for two days. After the culture was subjected to centrifugation at 20,000 x g for 30 minutes, the supernatant was admixed with the same volume of 10% trichloro acetic acid and stirred at 4°C for 16 hours. This solution was then subjected to centrifugation at 10,000 x g for 30 minutes. The thus obtained precipitate was dissolved in 4.5 ml of 40 mM Tris-HCl, pH 8.8, and subsequently neutralized with (about 0.5 ml) 2 M Tris-base. This protein solution was given to the company Toplab Gesellschaft für angewandte Biotechnologie mbH, Martinsried, Germany, for protein sequencing. At this company, the protein solution was electrophoretically separated in SDS polyacrylamide gel, the gel was stained with Coomassie Blue and the staining was subsequently removed by 10% acetic acid. For the enzymatic digestion of the protein, the protein bands were cut from the gel, pressed through a sieve and fragmented (pores 30 µm x 100 µm). The crushed gel was then washed with half concentrated incubation buffer (12.5 mM Tris, 0.5 mM EDTA pH 8.5) for 2 minutes. Subsequently, it was subjected to centrifugation, the buffer was removed and the gel was dried in the "Speedvac" for one hour (about 5% residual water, rubber-like). Subsequently, a solution of endoproteinase LysC in 400 µl 12.5 mM Tris/HCl, pH 8.5 (enzyme: protein = 1 : 10) and 0.1% of laurylmaltoside was prepared. 200 µl of this solution were added to the sample and incubated in the heat block shaker at 37°C overnight. In order to elute the peptide fragments, a one hour incubation with 1% TFA was carried out, twice, followed by centrifugation, and subsequently by elution with 10% formic acid, 20% isopropanol, 60% acetonitrile for 3 hours. The peptide fragments obtained were then separated from each other by HPLC (column Superspher 60 RP select B (Merck, Darmstadt) 2 mm x 125 mm; buffer A 0.1% trifluoro acetic acid, buffer B: 0.085% TFA in acetonitrile; flow rate: 0.2 ml/min; gradient: 5-60% in 60 min; detection at 206 nm. The peptide fragments obtained were then sequenced in an automatic sequencer Procise 492 (Applied Biosystems,

PE); the procedure being the stepwise Edman degradation in a modification according to Hunkapiller (Hunkapiller et al., Meth. Enzymol. 91 (1983), 399-413). Six different peptide sequences (see Seq. ID Nos. 5 to 9, Seq. ID No. 21) were identified which were designated lysC-66, lysC-67, lysC-82, lysC-83, lysC-88 and "N-terminus".

Preparation of a genomic DNA library from *Leuconostoc mesenteroides* NRRL B1355

Leuconostoc mesenteroides NRRL-B1355 (purchased from ATCC) was cultured in 100 ml YT medium (Sambrook et al, loc. cit.) additionally containing 2% (w/v) of glucose and 50 mM sodium phosphate buffer pH 7.0, at 28°C for 36 hours. After harvesting the cells by centrifugation, genomic DNA was isolated according to Ausubel et al. (Current Protocols in Molecular Biology, Volume 1, Greene and John Wiley & Sons (1994), USA).

100 µg of genomic DNA from *Leuconostoc mesenteroides* NRRL-B1355 were partially digested with 0.001 units of the restriction enzyme Sau3A for 30 minutes, subsequently extracted with phenol:chloroform:isoamyl alcohol (25:24:1) and precipitated with ethanol. 2.5 µg of the partially digested DNA obtained from *Leuconostoc mesenteroides* NRRL-B1355 were ligated with T4 DNA ligase in 1 µg of the BamHI-cut and dephosphorylated vector pBKCMVBamHI (Stratagene) under the conditions indicated by the manufacturer (Stratagene, pBK phagemid vectors instruction manual & T4 DNA ligase ligation kit). 2 µl of the ligation mixture were packaged with Gigapack III Gold (Stratagene) according to the instructions of the manufacturer and stored after the amount of phage content had been determined.

Preparation of the probe for isolating the alternansucrase gene

From the peptide sequences lysC-66 (Seq. ID No. 5), lysC-67 (Seq. ID No. 6), lysC-82 (Seq. ID No. 7), lysC-83 (Seq. ID No. 8) and lysC-88 (Seq. ID No. 9) obtained after tryptic digestion of the purified alternansucrase protein (see above) the peptides lysC-82 and lysC-83, after having undergone reverse translation, were selected for the synthesis of degenerated oligonucleotides (Seq. ID No. 10, Seq. ID No. 11). Said oligonucleotides served as primers in a PCR reaction on genomic

DNA of NRRL-B1355. All positions within oligonucleotides depicted as N were replaced by inosin in the primer synthesis.

PCR Reaction conditions

The reaction mixture was prepared with the buffers supplied for Taq polymerase (Company GibcoBRL).

Reaction mixture:

Taq Polymerase (Gibco)

DNA	100	ng (genomic NRRL-B1355)
DNTPs	2.5	mM for each nucleotide
primer	10	μl of a solution containing 0.2 μMol
10 fold buffer	5	μl
magnesium chloride	2	mM
polymerase	1	unit
water	ad	50 μl

Step 1	95°C	3'
Step 2	95°C	1'
Step 3	58°C	2'
Step 4	72°C	2'
Step 5	72°C	10'

40 repetitions of steps 2 to 4

An 837 bp fragment (Seq. ID No. 12) resulting from this PCR reaction, the ends of which were blunted with T4 DNA polymerase, was cloned into the SmaI-cut pBlueSkript vector (Stratagene). The resulting plasmid was designated pAlsu-PCR-lysc82/83. After sequencing of the insert and computer-aided translation into the corresponding protein sequences, a data base comparison was carried out in the Swiss Prot data base. This comparison showed homologies to known glycosyl transferases (P49331, P11001, P68987, P13470, P27470, P29336).

About 5,000 phages of the genomic DNA library of *Leuconostoc mesenteroides* NRRL-B1355 were plated out using the bacterial strains and nutrient solutions indicated by the manufacturer (Stratagene), and after incubation at 37°C for 12 hours were transferred to nitrocellulose filters. This was followed by denaturation of the phages by immersion of the nitrocellulose filters in 1.5 M sodium chloride, 0.5 M caustic soda solution for 2 minutes and neutralization of the filters by immersion in 1.5 M sodium chloride, 0.5 M Tris-HCl, pH 8.0 for 5 minutes. After rinsing the filters in 0.2 M Tris-HCl, 2 x SSC, the phage DNA was bound to the membranes by UV cross link (Stratalinker of the company Stratagene, 120,000 μ J for 30 seconds). The filters were incubated in a prehybridization solution (5 x SSC, 0.5% BSA, 5 x Denhardt, 1% SDS, 40 mM sodium phosphate buffer, pH 7.2, 100 mg/l herring sperm-DNA, 25% formamide) at 42°C for 6 hours. 30 ng of the isolated insert from the plasmid pAlsu-PCR-lysc82/83 were radioactively labeled by means of a multiprime kit (Boehringer Mannheim) using α -³²P dCTP (ICN Biomedicals). This radioactive probe was added to the prehybridization mixture and the filters were incubated in this hybridization mixture at 42°C overnight. After removal of the hybridization mixture the filters were washed three times in a washing solution (0.1 x SSC, 0.5% SDS) at 55°C for 15 minutes. An X-ray film (Kodak) was then placed on the filter for 18 hours. Phage colonies, producing hybridization signals, were identified, isolated, resuspended in SM medium and then again plated out in a dilution such that they could be recognized as single plaques. After these phages were transferred to nitrocellulose filters and subjected to further treatment and hybridization under conditions as described above, hybridizing phages were obtained as individual isolates by means of the radioactive gene probe used. After *in vivo* excision of the isolated phages in accordance with the manufacturer's instructions (Stratagene) the clones AS-19B1 and AS-19B2 could be isolated as plasmids. After complete sequencing of both clones (Agowa) (Seq. ID No. 13, Seq. ID No. 14) both sequences showed an 1008 bp overlap. The joining of Seq. ID No. 13 with Seq. No. 14 followed by computer aided translation of all possible reading frames allowed a continuous reading frame, starting with the codon ATG (corresponding to the bases 678 to 680 in Seq. ID No. 1), to be identified. As no stop codon could be found in this composed reading frame, additional clones were isolated in order to obtain the complete coding sequence of alternansucrase.

Therefore, about 5,000 phages of the genomic DNA library of *L. mesenteroides* NRRL-B1355 were again examined for hybridization by means of a clone AS-19B2 subfragment radioactively labeled using the multiprime kit (Boehringer Mannheim), as described above. The hybridization probe was prepared with the use of the HindIII (restriction site in the insert of AS-19B2) / Sall (cuts the pBKCMV phagemid vector in the polylinker)-fragment from AS-19B2. Said fragment contains 372 bases of the 3' end of the sequences encoding the above-described reading frame. The screening of the phage library, singling out, and transformation of the phages into plasmids was carried out under the above-described conditions. After complete sequence analysis of the thus isolated clones AS-28B (see Seq. ID No. 15) and AS-29Ba (Seq. ID No. 16) it was possible to identify an overlap of 960 identical bases (corresponding to bases 4863 to 5823 in Seq. ID No. 1) between clones AS-19B2 (Seq. ID No. 14) and AS-28B and an overlap of 567 identical bases (corresponding to bases 5256 to 5823 in Seq. ID No. 1) between clones AS-19B2 and AS-29Ba (Seq. ID No. 16). Clones AS-28B and AS-29Ba have 1523 identical bases (corresponding to bases 5256 to 6779 in Seq. ID No. 1). After computer-aided joining of clones AS-19B1, AS-19B2 and AS-28B a continuous reading frame starting with codon ATG (bases 678 to 680 on the complete sequence) appeared. This reading frame also does not contain a stop codon. After the joining of clones AS-19B1, AS-19B2, AS-28B and AS-29Ba it was possible to identify a reading frame starting with the codon "ATG" (corresponding to bases 678 to 680 in Seq. ID No. 1) and ending with "TAA" (corresponding to bases 6849 to 6851 in Seq. ID No. 1) encoding 2057 amino acids. In addition to the coding region, the entire isolated and identified DNA sequence of the composed clones (Seq. ID Nos. 13 - 16) contains 677 bases in the 5' region and 2469 bases in the 3' region which represent sequences not encoding alternansucrase (see Fig. 1).

Example 2: Construction of plasmid pAlsu-pSK for the transformation of *E. coli* and test of the protein extracts for enzymatic activity

Plasmids AS-19B1, AS-19B2, AS-28B and AS-29Ba (see Example 1) were joined in the following manner: A NotI-(restriction site in the polylinker of vector pBK CMV, company Novagen)/Clal-fragment of clone AS-19B1 was inserted into the vector

pBluescript SK (company Stratagene) at the same restriction sites (= first cloning step). Consecutive insertion of the *Clal/XhoI* fragment from AS-19B2, *XhoI/MluI* fragment from AS-28B and *MluI/BsaBI* (*BsaBI*-cut fragment cloned into the blunted *Apal* restriction site of the vector) fragment of AS-28B into the clone obtained from the first cloning step produced plasmid pAlsu-pSK (see Fig. 2). This plasmid contains the complete coding sequence of the alternansucrase from *Leuconostoc mesenteroides* NRRL-B1355 as well as non-coding sequences of 677 bp (promoter region) in the 5' region and 539 bp in the 3' region (Seq. ID No. 17).

Plasmid pAlsu-pSK was then transformed in *E. coli* (DH5 α company Fermentas). The bacteria were then cultured at 27°C for two days in 50 ml "Terrific broth" (the composition of which is described in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (supplemented with 0.5% glucose) or in a fermentation medium having the following composition: KH_2PO_4 1.5 g/l, $(\text{NH}_4)_2\text{SO}_4$ 5.0 g/l, NaCl 0.5 g/l, Na-citrate 1.0 g/l, $\text{Fe}^{2+}\text{SO}_4 \times 7 \text{H}_2\text{O}$ 0.075 g/l, yeast extract 0.5 g/l, tryptone 1.0 g/l, glucose 15.0 g/l, $\text{MgSO}_4 \times 7 \text{H}_2\text{O}$ 0.3 g/l, $\text{CaCl}_2 \times 2 \text{H}_2\text{O}$ 0.014 g/l, mineral salts 10 ml/l, H_3BO_3 2.5 g/l, $\text{CoCl}_2 \times 6 \text{H}_2\text{O}$ 0.7 g/l, $\text{CuSO}_4 \times 5 \text{H}_2\text{O}$ 0.25 g/l, $\text{MnCl}_2 \times 4 \text{H}_2\text{O}$ 1.6 g/l, $\text{ZnSO}_4 \times 7 \text{H}_2\text{O}$ 0.3 g/l, $\text{Na}_2\text{MoO}_4 \times 2 \text{H}_2\text{O}$ 0.15 g/l, vitamin B1 (thiamine) 0.005 g/l.

All cultures contained 100 mg/l ampicillin. The cells were then harvested by centrifugation, resuspended in 2 ml 50 mM Na-phosphate buffer pH 7.2 and crushed by a French Press. Subsequently, they were again subjected to centrifugation to remove solid particles of the crushed cells, and the supernatant (hereinafter referred to as (protein) extract) was used after sterilfiltration (Sterivex GV 0.2 μm , millipore) for further analyses.

In vitro preparation of alternan by means of protein extracts

For the in vitro preparation of alternan, 200 μl each of the extracts obtained were examined in 2 ml each of 100 mM Na-citrate buffer pH 6.5 and 20% (w/v) saccharose for activity in the presence and absence of 100 μl of 10 mM maltose. The reaction mixture was incubated at 37°C for 24 hours. In the subsequent

precipitation with the same volume of ethanol in the absence of maltose no precipitable polymer was found. In the batch containing maltose, HPLC chromatography (Dionex PA-100 column, running buffer 150 mM NaOH, elution buffer 150 mM NaOH + 3 M sodium acetate buffer gradient) showed the formation of oligomers (see Fig. 3).

Activity gel

20 ml each of the individual protein extracts were applied to a 6% SDS-PAA gel and separated at a current strength of 20 mA per gel. (Before application to the gels, the extracts were not incubated at 95°C). Subsequently, the extracts were examined for sucrase activity according to the method of Miller and Robyt (Analytical Biochemistry 156 (1986), 357-363).

The control (dextransucrase NRRL-B-512F, see Example 3 for its preparation) showed polymerizing activity. The protein extracts of the above-described *E. coli* cells containing the plasmid pAlsu-pSK, did not show any polymer-forming activity.

Example 3: Cloning and expression of dextransucrase from *Leuconostoc mesenteroides* NRRL-B512F

Isolation of genomic DNA

Leuconostoc mesenteroides NRRL-B512F (obtained from ATCC) was cultured at 28°C for 48 hours in YT-medium (Sambrook et al., Molecular Cloning: A Laboratory Course Manual, 2nd edition (1989), Cold Spring Harbor Press, New York) additionally containing 1% of saccharose and 50 mM sodium phosphate buffer pH 7.0. After harvesting the cells by centrifugation, genomic DNA was isolated according to Ausubel et al. (Current Protocols in Molecular Biology, Volume 1, Greene and John Wiley & Sons (1994), USA).

PCR Amplification of the Dextransucrase Gene and Cloning in pET24a

For the recombinant expression of dextransucrase in *E. coli*, the gene encoding dextransucrase was cloned in the expression vector pET24a (Novagen) after PCR amplification. For this purpose, an EagI restriction site was introduced at the 5' end of the sequences encoding the dextransucrase and an XhoI restriction site at the 3'

end, together with the PCR primers used (5'b512-1: 5'-ACTgCggCCgCATgCCATTTACAgAAAAg-3'; Seq. ID No. 3 and 3'b512: 5'-ACTgCTCgAgTTATgCTgACACAgCATTTC-3'; Seq. ID No. 4) derived from the sequence of WO 89/12386. Subsequent cloning into the corresponding restriction sites of the polylinker of pET24a was carried out. The resulting plasmid was designated UL5-20.

PCR Reaction conditions

Buffer and polymerase of the company Gibco BRL were used.

DNA:	100	ng (genomic NRRL-B512F)
10 fold buffer	5	μl
MgCl ₂	4	mM
5' primer	50	ng
3' primer	50	ng
dNTP	1	mM of each nucleotide
Pfu polymerase	0.5	units
water	ad 50	μl

step 1	95°C	4 minutes
step 2	95°C	1 minute
step 3	55°C	1 minute
step 4	72°C	5 minutes
step 5	72°C	10 minutes

40 repetitions were made between steps 2 and 4.

Preparation of recombinant dextranucrase

BL21(DE3) *E. coli* cells containing the plasmid UL5-20 were cultured in YT medium (see above) at 37°C up to an OD₆₀₀ = 0.8. Subsequently, the cells were subjected to induction with 0.2 mM IPTG and cultured anew at 18°C for 24 hours. After harvesting the cells by centrifugation and resuspending them in sodium phosphate buffer, pH 5.2, the cells were crushed in a French Press. The solution obtained

was freed from insoluble components by centrifugation and the supernatant containing dextranucrase and referred to hereinafter as the extract was obtained.

Example 4: PCR Amplification of the Coding Region of Alternansucrase and Cloning in pET24a

The coding region of alternansucrase was amplified in a PCR reaction (see the reaction conditions below) with genomic DNA from the *Leuconostoc mesenteroides* strain NRRL-B1355 as a template. An NheI restriction site was introduced at the 5' end by means of primers A1-4 (Seq. ID No. 18), and a Sall-restriction site at the 3' end by means of primer A1-5 (Seq. ID No. 19). A fragment of about 6200 bp was isolated.

A1-4: 5'-GGG CCC GCT AGC ATG AAA CAA CAA GAA ACA GT

A1-5: 5'-CCC GGG GTC GAC CTT TGT CGA ATC CTT CCC

Reaction conditions of the PCR (kit of the company Gibco BRL):

DNA	1 µl
10 x buffer	5 µl
10 mM per dNTP	2 µl
50 mM MgSO ₄	2 µl
primer per	1 µl
Platinum DNA polymerase	0.2 µl
distilled water	37.8 µl
step 1	95°C, 2 minutes
step 2	95°C, 20 seconds
step 3	47°C, 20 seconds
step 4	68°C, 7 minutes (prolonged by 3 seconds per cycle)
step 5	68°C, 15 minutes

Steps 2 to 4 were repeated 35 times altogether before step 5 was carried out.

The PCR fragment obtained was purified according to standard methods, treated with the restriction endonucleases NheI and Sall, ligated into vector pET24a (of the company Novagen) which had likewise been cut with these enzymes, and the ligation product was transformed into *E. coli*. After preparation of the plasmid and restriction digestion, three positive clones were selected. They were designated pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21 (see Fig. 4), respectively. All contained the sequence indicated in Seq. ID No. 20 as an insertion.

Example 5: Expression of the Recombinant Alternansucrase in *E. coli* in Shake Flask Cultures and in the Fermenter

Shake Flask Culture

Plasmids pAlsu-pET24a-3, pAlsu-pET24a-7, pAlsu-pET24a-21 and pET24a were transformed into *E. coli* BL21 (DE3), of the company Novagen, and after initial culturing at 37°C for 3 hours in 3 ml YT medium (Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd edition (1989), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY) they were each cultured in shake flasks in 2 replicas in 50 ml Davis minimal medium (DIFCO Manual, Dehydrated Culture Media and Reagents for Microbiology, 10th edition, Detroit Michigan, USA (1984)) containing 0.2% glucose instead of dextrose as a carbon source at 37°C until an OD₆₀₀ of about 0.8 was reached. After centrifugation and resuspension, one of the two replica cultures was cultured in Davis Minimal Medium (DMA) containing 1% lactose as the carbon source and inductor at 27°C for another 16 hours. The cells of the individual cultures were harvested after centrifugation, resuspended in 50 mM sodium acetate buffer pH 5.3, and a protein extract was prepared as described in Example 2.

Fermenter

Clone pAlsu-pET24a-21 transformed in *E. coli* BL21(DE3) was cultured in a 2 l fermenter (Biostad B; B.Braun, Melsungen) under the following conditions:

Medium:

Fermentation medium: KH_2PO_4 1.5 g/l, $(\text{NH}_4)_2\text{SO}_4$ 5.0 g/l, NaCl 0.5 g/l, Na-citrate 1.0 g/l, $\text{Fe}^{2+}\text{SO}_4 \times 7 \text{ H}_2\text{O}$ 0.075 g/l, yeast extract 0.5 g/l, tryptone 1.0 g/l, glucose 15.0 g/l, $\text{MgSO}_4 \times 7 \text{ H}_2\text{O}$ 0.3 g/l, $\text{CaCl}_2 \times 2 \text{ H}_2\text{O}$ 0.014 g/l, mineral salts 10 ml/l, H_3BO_3 2.5 g/l, $\text{CoCl}_2 \times 6 \text{ H}_2\text{O}$ 0.7 g/l, $\text{CuSO}_4 \times 5 \text{ H}_2\text{O}$ 0.25 g/l, $\text{MnCl}_2 \times 4 \text{ H}_2\text{O}$ 1.6 g/l, $\text{ZnSO}_4 \times 7 \text{ H}_2\text{O}$ 0.3 g/l, $\text{Na}_2\text{MoO}_4 \times 2 \text{ H}_2\text{O}$ 0.15 g/l, vitamin B1 (thiamine) 0.005 g/l.

Carbon source: Glucose (1.5% (w/v)) is present in the medium, 70% (w/v) glucose solution is added.

Automatic pH control by ammonia and phosphoric acid at pH 7.0 \pm 0.1. A 20% concentration of pO_2 is adjusted in the medium via control by the stirrer.

Conditions:

1.5 l of fermentation medium were inoculated with 50 ml of the preculture.

The cells were first cultured at 37°C until the glucose present was consumed. They were then cultured at the same temperature at a feeding rate of 9 g of glucose $\times \text{l}^{-1} \times \text{h}^{-1}$ until an $\text{OD}_{600} = 40$ was reached. At this time, the temperature of the culture broth was lowered to 20°C and the amount of glucose addition was lowered to 2 g $\times \text{l}^{-2} \times \text{h}^{-1}$. At a culture temperature of 20°C, the culture was subjected to induction with 0.2 mM IPTG (isopropyl- β -D-thiogalactopyranoside (Sigma)). After culturing at 20°C for another 18 hours, the cells were harvested by centrifugation, resuspended in 50 mM sodium phosphate buffer pH 5.3 and an extract was prepared as described in Example 2.

Example 6: SDS PAGE Assay of the Activity of the Recombinant Alternansucrase, Periodic Acid Oxidation and Staining according to Schiff

Protein extracts were prepared from *E. coli* shake flask cultures (strain BL21 (DE3)), containing the plasmids pAlsu-pET24a-3, pAlsu-pET24a-7, pAlsu-pET24a-21 and pET24a (control), respectively. Two different extracts were each prepared from the cells transformed with the different extracts, one of said extracts being prepared before induction with IPTG and the other one being prepared after induction with IPTG at the end of culturing. The activity of these extracts of shake

flask cultures (see Example 5) was detected by SDS PAGE separation of the proteins, followed by SDS removal by washing with 50 mM sodium acetate buffer pH 5.3 and incubation of the gels in 50 mM sodium acetate pH 5.3, 5% (w/v) saccharose at 37°C for 16 hours, followed by periodic acid oxidation of the polymer formed and staining by means of acidic Schiff reagent (Miller and Robyt, Analytical Biochemistry 156, (1986), 357-363).

Fig. 5 shows that sucrase activity has not been found for either one of the extracts (preparation of the extract before and after IPTG-induction) containing the cloning vector pET24a. In the case of strains which had been transformed with the plasmids pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21, respectively, all protein extracts showed sucrase activity at the end of the induction phase (concentrated in one band).

Before induction with IPTG such activity bands were not found.

As the polymer formed in the gel can be stained according to the above-described methods by acidic Schiff reagent, it can be assumed not to be composed of pure α -1,3-linked units which would not lead to any staining.

As the gene contained in vectors pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21, respectively, was isolated from the *Leuconostoc mesenteroides* strain NRRL-B1355 which expresses at least one dextran sucrase apart from alternansucrase, it was not possible to determine unambiguously with this staining method whether the nucleic acid sequence contained in the plasmid actually encodes an alternansucrase. Dextrans and alternans can both be detected by this method because both polymers contain α -1,6 linkages.

Example 7: Tests for the Enzymatic Activity of Recombinantly Prepared Alternansucrases after Heat Treatment and for the Specificity of Alternansucrase

In order to prove polymerization activities, extracts from shake flask cultures were used (see Example 5). 100 μ l of extract were each added to 2 ml reaction buffer (50 mM sodium acetate pH 5.3, 20% saccharose) and incubated at 37°C for 24 hours. For comparison, an extract inactivated by a 10 minute treatment at 95°C, and an

extract from *E. coli* BL21(DE3) containing vector pET24a were used. Polymer formation was only found in the batch that had not been inactivated, while the batch treated at 95°C for 10 minutes and the batch with the extract from BL21(DE3) containing pET24a did not show any polymer formation. After addition of the same volume of absolute ethanol to all batches, polymers could only be precipitated from the batch which had not been inactivated. This finding is a clear indication of the activity of alternansucrase, because the dextransucrase present in NRRL B-1355 is inactivated by a treatment at 45°C for 30 minutes, while alternansucrase remains active under these conditions (Lopez-Munguia et al., Enzyme Microb. Technol. 15 (1993), 77-85). The enzymatic assay by a coupled enzymatic test of the glucose and fructose released and of the saccharose still contained in the reaction mixture after 24 hours, respectively, revealed that fructose was only present in the extract that was not inactivated.

For carrying out the enzymatic test either purified protein or crude protein extract is added in different dilutions to 1 ml batches containing 5% saccharose and 50 mM acetate, pH 5.5 and subjected to incubation at 37°C. After 5 minutes, 10 minutes, 15 minutes, 20 minutes, 25 minutes and 30 minutes, 10 µl each are removed from these batches and the enzymatic activity of alternansucrase is terminated by immediate heating to 95°C. Subsequently, in the coupled photometric test, the portions of fructose and glucose released by alternansucrase and the portion of used-up saccharose, respectively, are determined. For this purpose, 1 µl to 10 µl of the inactivated sample are placed into 1 ml of 50 mM imidazole buffer, pH 6.9, 2 mM MgCl₂, 1 mM ATP, 0.4 mM NAD and 0.5 U/ml hexokinase. After sequential addition of about 1 u of glucose-6-phosphate dehydrogenase (from *Leuconostoc mesenteroides*), about 1 u of phosphoglucose isomerase and about 5 u of invertase, the alteration of adsorption at 340 nm is measured. Subsequently, the amount of fructose and glucose released and used-up saccharose, respectively, is calculated according to the Lambert-Beer law.

In control batches (inactivation of the extract by treatment with 95°C and extract from *E. coli* containing pET24a) no significant release of fructose and no decrease of saccharose, respectively, was found in the reaction batch after 24 hours.

These results confirm that the specificity of the sucrase encoded by plasmids pAlsu-pET24a-3, pAlsu-pET24a-7 and pAlsu-pET24a-21, respectively, is that of a

glucosyltransferase. The specificity of a fructosyl transferase, the presence of which has been described for some strains of the genus *Leuconostoc* is to be excluded, because otherwise glucose should have been found.

Example 8: Production of Alternan by Means of Alternansucrase Prepared in *E. coli*

100 ml of extract obtained by fermentation of *E. coli* BL21(DE3) containing plasmid pAlsu-pET24a-3 (see Example 4) were added to 900 ml of reaction buffer (50 mM sodium acetate pH 5.3, 20% saccharose) and incubated at 37°C for 24 hours. The addition of the same amount of absolute ethanol to the reaction mixture caused the alternan formed to precipitate. After the precipitate was washed twice with 50% ethanol, it was dried by lyophilization. The yield of dried polymer based on the amount of saccharose used in the reaction was 60%.

Example 9: HPLC Analysis of Alternan and Dextran After Dextranase Digestion

100 mg of the polymer prepared in Example 7 and 100 mg of dextran T10 (Pharmacia) were each dissolved in 1 ml of water. 40 µl each of these solutions were added to 700 µl reaction buffer (50 mM potassium phosphate pH 5.7, 8 units of dextranase, ICN Biomedicals Inc. No. 190097), and incubated at 37°C for 16 hours. 50 µl of the polymer solutions not treated with dextranase (see Fig. 6) and 50 µl of the polymer solutions treated with dextranase (Fig. 7) were analyzed by HPLC (Dionex, column PA-100, NaOH / NaOH-NaAc gradient).

In the case of dextran T10 the cleavage of the polymer into different molecules of lower molecular weights can be clearly seen. The entire high molecular weight dextran is converted by dextranase into smaller units (mostly isomaltose). By contrast, in the case of alternan, short chained oligosaccharides only appear in small amounts after dextranase incubation. Most of the alternan is not digestible by dextranase. This finding suggests that the product prepared by recombinant alternansucrase is not dextran, but alternan which is known to be hardly accessible

to enzymatic digestion by dextranase (Lopez-Mungia et al., Enzyme Microb. Technol. 15, (1993), 77-85).

Example 10: *In vitro* Preparation of Alternan in the Absence of Dextranase

100 μ l extract from shake flask cultures (see Example 5) were added to 2 ml of reaction buffer (50 mM sodium acetate, pH 5.3, 20% saccharose). 50 units of dextranase (Biomedicals Inc. No. 190097) were additionally added to another batch. Two corresponding batches which contained dextransucrase from *Leuconostoc mesenteroides* NRRL-B512F instead of the enzyme extract served as controls; one of these two batches had dextranase additionally admixed to it.

After precipitation with ethanol, the reaction batch with dextransucrase and dextranase did not show any polymer formation. All other batches were found to show polymer formation.

Example 11: *In vitro* preparation of Oligoalternan and HPLC Analysis

Oligoalternan was prepared as described in Example 2, with a protein extract in the presence of maltose and was subsequently detected (See Fig. 8) by HPLC-chromatography (see Example 2). For comparison, a portion of this batch was admixed with 50 units of dextranase (Biomedicals Inc. 190097) after preparation of oligoalternan and subsequently separation by HPLC chromatography was carried out as well (see Fig. 9). A comparison of the two chromatograms shows that not only the height of the two peaks which can be allocated to the oligoalternan (α and β -anomer) (retention time between 15.87 and 16.61 minutes) but also the height of all the other peaks, the first signs of which are already visible without dextranase, remain unchanged. This finding suggests that recombinantly prepared alternansucrase allows oligoalternan to be prepared without the simultaneous production of oligodextran. Oligodextran would be liable to digestion by dextranase, which would have to show up in a decrease of the height of the peaks in the HPLC chromatogram, if oligodextran were present.

Example 12: Methylation analysis of Alternan

In order to further analyze the alternan produced *in vitro* a methylation analysis was carried out:

Permethylation

The permethylation was performed as described by Ciucanu and Kerek (Carbohydr. Res. 131 (1984), 209-218) by using NaOH/Mel in DMSO or by using a modified method according to Hakomori (Journal of Biochemistry 55 (1964 FEB), 205-208) which relies on the use of freshly prepared Li-Dimsyl/Mel (Dimsyl= methylsulfinyl carbanion) in DMSO at room temperature.

All reactions are performed under a nitrogen atmosphere. The permethylation products are isolated by extracting the excess of methyl iodide by the use of dichloromethane. DMSO and salts were washed out at the end.

Degradation into partially methylated sorbitacetates (methylation analysis)

The permethylated glucans were hydrolyzed with 2N trifluoroacetic acid at 120°C for 1-3 hours. After cooling the acid was removed by nitrogen. Then the resulting glucans were co-distilled with a small amount of toluene, afterwards reduced by NaBD₄ in 1N ammonia and finally, acetylated by pyridine/acetic anhydride (3h, 90°C). The products were extracted by dichloromethane and washed with NaHCO₃. The products in the organic phase were analyzed by gas chromatography.

Analysis of the acetylated products

The acetylated products were analyzed by gas chromatography which was performed with a chromatograph manufactured by the Carlo-Erba company model GC 6000 Vega equipped with an on-column injector, a 25m CPSol8CB and a FID-detector. As a carrier gas hydrogen (80 kPa) was used.

The identification and integration of the peaks was performed as described by Sweet et al. (Carbohydr. Res. 40 (1975), 217).

Results

The following main components were identified by gas chromatography:

Sorbit acetylated in position	Interpretation
1,5	Terminal Glucopyranose
1, 3, 5	3-linked Glucopyranose
1, 5, 6	6-linked Glucopyranose
1, 3, 5, 6	3,6-linked Glucopyranose

Furthermore, small amounts (rel. amount 0.2-0.4 mol%) of the following components were also found: 1, 4, 5- and 1, 3, 4, 5-sorbit and another tetraacetyl component (1,5,x,y). It is supposed that these components are due to incomplete methylation.

The following amounts were found for the above mentioned components in different experiments which were performed by changing the length of hydrolysis (indicated in bold by the number of hours) (MA=methylation analysis1; MA-b= methylation analysis 2):

Values in mol%

Ac in Pos	MA (1h)	MA (2h)	MA (3h)	MA-b(2h)
1,5	10,49	10,56	9,17	12,71
1,3,5	31,69	34,70	32,95	23,12
1,4,5	0,70	0,30	0,36	0,33
1,5,6	47,02	44,17	47,23	54,62
1,3,4,5	0,27	0,22	0,25	0,31
1,5,x,y	0,19	0,32	0,36	0,24
1,3,5,6	9,64	9,73	9,68	8,67

Example 13: Construction of an expression cassette for plants: vacuolar and plastidic expression of an alternansucrase.

By using plasmid Alsu-pET24a as a template and the PCR primers Al-5'-1.2 and Al-3'-2.2 (see SEQ ID NO 53 and 54) we amplified the coding region of alternansucrase from *Leuconostoc mesenteroides* which was then cut by the

restriction enzymes Sall and PstI. Afterwards the resulting fragments were cloned into Sall and Sdal digested plasmids a) pBinAR-pat-Hyg and b) pBinAR-fnr-Hyg. The resulting plasmids were called a) pat-Alsu-Hyg (see figure 11) and b) fnr-Alsu-Hyg (see figure 12).

Note: The bacterial secretion signal peptide was removed from the cds by choice of the PCR primers.

PCR conditions:

Buffer and polymerase from Boehringer Mannheim (Pwo Polymerase No. 1644947)

DNA	0,5 ng
10x Buffer + MgSO ₄	5 µl
dNTPs (je 10 mM)	2 µl
Primer Sp-AS-5'	100 nM
Primer Sp-AS-3'	100 nM
Pwo Polymerase	1,0 unit
distilled water	ad 50 µl

Reaction conditions:

Step 1	95°C	2:30 min
Step 2	95°C	0:30 min
Step 3	47°C	0:30 min
Step 4	68°C	7:00 min
	(plus 3 sec per cycle)	
Step 5	68°C	15:00 min

The steps 2 to 4 were repeated 35 times in a cyclical manner.

Example 14: Northern blot analysis for expression of alternansucrase in transgenic plants

Leaves or tubers from potato plants transformed via agrobacteria with plasmids pat-Alsu-Hyg and fnr-Alsu-Hyg, respectively, were pulverized in a mill, type MM 200, (Retsch GmbH & Co. KG, 42781 Haan, Germany) at 30 Hz for 50 sec. RNA was extracted according to Logemann et al. (Anal. Biochem. 163 (1987), 16-20). 50 µg

RNA per sample were loaded on 1% agarose gels containing formaldehyde. After electrophoresis the RNA was transferred to nylon membranes (Hybond N, Amersham, UK) by the capillary transfer method (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)). Fixation of nucleic acids at the membrane was achieved by UV crosslinking (Stratalinker by Stratagene).

Membranes were prehybridized at 42°C in hybridization buffer (25% (v/v) formamide, 250 mM sodium phosphate, pH 7.2, 250 mM sodiumchloride, 1 mM EDTA 7% (w/v) SDS, 25% (w/v) polyethyleneglycol 6000, 0,25 mg/ml sheared salmon sperm DNA) for 6 h. Afterwards hybridization was performed at 42°C over night in hybridization buffer containing a radiolabelled probe in addition. The radioactive probe was prepared by using the Random Primed DNA Labelling Kit (Boehringer Mannheim, 1004760) and the approx. 4 kb KpnI/XhoI-fragment from plasmid pAlsu-pSK according to the manufacturers manual. Membranes were washed at 50°C once for 20 min in 3xSSC (Sambrook et al., Molecular cloning: A laboratory manual, 2nd issue; Cold Spring Harbor Laboratory Press, NY, USA (1989)) followed by washing once for 20 min in 0.5xSSC before exposing the membrane to an x-ray-film over night.

Claims

1. A nucleic acid molecule encoding a protein having the biological activity of an alternansucrase, selected from the group consisting of
 - (a) nucleic acid molecules encoding at least the mature form of a protein comprising the amino acid sequence indicated in Seq. ID No. 2 or the amino acid sequence which is encoded by the cDNA contained in plasmid DSM 12666;
 - (b) nucleic acid molecules comprising the nucleotide sequence indicated in Seq. ID No. 1 or the nucleotide sequence of the cDNA contained in plasmid DSM 12666 or a corresponding ribonucleotide sequence;
 - (c) nucleic acid molecules encoding a protein, the amino acid sequence of which has a homology of at least 40% to the amino acid sequence indicated in Seq. ID No 2;
 - (d) nucleic acid molecules, one strand of which hybridizes with any one of the nucleic acid molecules as defined in (a) or (b);
 - (e) nucleic acid molecules comprising a nucleotide sequence encoding a biologically active fragment of the protein which is encoded by any one of the nucleic acid molecules as defined in (a), (b), (c) or (d); and
 - (f) nucleic acid molecules, the nucleotide sequence of which deviates on account of the degeneration of the genetic code from the sequence of the nucleic acid molecules as defined in (a), (b), (c), (d) or (e).
2. An oligonucleotide or polynucleotide which specifically hybridizes with a nucleic acid molecule of claim 1.
3. A vector containing a nucleic acid molecule according to claim 1.
4. The vector according to claim 3, wherein the nucleic acid molecule is connected in sense orientation to regulatory elements ensuring the transcription and synthesis of a translatable RNA in prokaryotic or eukaryotic cells.
5. Plasmid pAlsu-pSK deposited under the accession No. DSM 12666.

6. A host cell transformed with a nucleic acid molecule of claim 1 or a vector of claim 3 or 4 or descended from such a cell.
7. The host cell according to claim 6, which is a cell of a microorganism.
8. The host cell according to claim 6, which is an *E. coli* cell.
9. A method for preparing a protein having the biological activity of an alternansucrase or a biologically active fragment thereof, wherein a host cell of any one of claims 6 to 8 is cultured under conditions permitting the synthesis of the protein, and wherein the protein is isolated from the cultured cells and/or the culture medium.
10. A protein or biologically active fragment thereof encoded by a nucleic acid molecule of claim 1 or prepared according to the method of claim 9.
11. A transgenic plant cell transformed with a nucleic acid molecule of claim 1 or a vector of claim 3 or 4, or descended from such a cell, wherein said nucleic acid molecule encoding the protein having the biological activity of an alternansucrase is under the control of regulatory elements permitting the transcription of a translatable mRNA in plant cells.
12. A plant containing the plant cells of claim 11.
13. The plant according to claim 12, which is a useful plant.
14. The plant according to claim 12 or 13, which is a sugar-storing or starch-storing plant.
15. Propagation material of a plant according to any one of claims 12 to 14, containing the plant cells of claim 11.

16. A method for preparing alternan comprising the steps of extracting and isolating the alternan from a plant according to any one of claims 12 to 14.
17. Alternan obtainable from a plant according to any one of claims 12 to 14, from propagation material according to claim 15 or by a method according to claim 16.
18. A method for preparing alternan and/or fructose, wherein
 - (a) a host cell of any one of claim 6 to 8 secretes an alternansucrase into a saccharose-containing culture medium; and
 - (b) alternan and/or fructose is/are isolated from the culture medium.
19. The method according to claim 18, wherein the host cell is immobilized.
20. A method for preparing alternan and/or fructose, wherein
 - (a) a saccharose-containing solution is contacted with a protein of claim 10 under conditions permitting the conversion of saccharose into alternan and/or fructose; and
 - (b) alternan and/or fructose is/are isolated from the solution.
21. The method according to claim 20, wherein the protein is immobilized on a carrier material.
22. A method for preparing alternan and/or fructose, wherein
 - (a) a saccharose-containing solution is contacted with a protein of claim 10 and acceptor molecules under conditions permitting the conversion of saccharose into alternan and/or fructose; and
 - (b) alternan and/or fructose is/are isolated from the solution.
23. The method according to claim 22, wherein the acceptor molecule is selected from the group consisting of maltose, isomaltose, isomaltotriose and methyl- α -D-glucan.

24. The method according to claim 22 or 23, wherein the protein is immobilized.
25. Cosmetic products or food products containing alternan as specified in claim 17 or prepared according to any one of claims 16 and 18 to 24.
26. A method for preparing cosmetic products or food products comprising a process according to any of claims 16 and 18 to 24 and the formulation of the resulting alternan in a form suitable for use as a cosmetic product or foodstuff.

1/12




Fig. 1

2/12




Fig. 2

3/12




Fig. 3

4/12




Fig. 4




Fig. 5

6/12




Fig. 6

7/12




Fig. 7

8/12




Fig. 8

9/12




Fig. 9

10/12




Fig. 10

11/12





Fig. 11

**Fig. 12**

SEQUENCE LISTING

<110> MAX-PLANCK-GESELLSCHAFT ZUR FÖRDERUNG DER WISSENSCHAFTEN e.V.

<120> Nucleic Acid Molecules Encoding Alternansucrase

<130> D 1042 PCT

<140>

<141>

<160> 54

<170> PatentIn Ver. 2.1

<210> 1

<211> 9321

<212> DNA

<213> Leuconostoc mesenteroides

<220>

<221> CDS

<222> (678)..(6848)

<400> 1

gatcaaatag atgctaccgt tatacgagac gttaaggatt atcccaatct agtcgttttg 60

cgtaccttgt cgaaagctta tggctctggca aacctgagaa tcggctatgg cgtcatgcag 120

gaaccacttt atcagggttat gcaggcogta cgtttaccat ataatttaaa tacctatcaa 180

atcacagggtg cagtagctgc ccttagtgat caactttatc tgcaatcagt tgttgctaag 240

gtgaagtctg aacgtgaaaa atttgaacaa tttttgacga aacaccagtt taagtattat 300

caatcacaaa ccaactttct ctggattaaa gttggtgatg cgaaacgtgt tggtgagggt 360

cttctgtcag aagggtatca aattaatgac cgcctaaatg ccgaatggat tcgcattgca 420

ttaggaactg tgtctgataa tgaggggatg cagcgcattt tattgaattg ttaaaaaata 480

gctaagagag tatgttcttc tcttacctat ttttatattg aattcctatt atttaatttt 540

gcatgacaat attaatagcg tgttacgatt ctactattta atgttaataa aattaataaa 600

tatggtatta tcttatatgg gtgatagatg caccaaatac tgtatcatgt ctggtcacat 660

gaaagggaga ataatta atg aaa caa caa gaa aca gtt acc cgt aaa aaa 710

Met Lys Gln Gln Glu Thr Val Thr Arg Lys Lys

1 5 10

ctt tat aaa tcc ggt aag gtt tgg gtt gca gca gct act gca ttt gcg 758

Leu Tyr Lys Ser Gly Lys Val Trp Val Ala Ala Ala Thr Ala Phe Ala

15 20 25

gta ttg ggg gtt tca act gta aca aca gtc cat gcg gat aca aat tcg 806

Val Leu Gly Val Ser Thr Val Thr Thr Val His Ala Asp Thr Asn Ser

30 35 40

aat gtc gct gtt aag caa ata aat aat aca gga acc aat gat tct ggc 854

Asn	Val	Ala	Val	Lys	Gln	Ile	Asn	Asn	Thr	Gly	Thr	Asn	Asp	Ser	Gly	
45						50					55					
gaa	aaa	aag	gta	ccg	gtt	cca	tca	act	aat	aat	gat	agt	ttg	aag	caa	902
Glu	Lys	Lys	Val	Pro	Val	Pro	Ser	Thr	Asn	Asn	Asp	Ser	Leu	Lys	Gln	
60					65					70					75	
gga	aca	gat	ggt	ttt	tgg	tat	gat	tca	gac	ggc	aat	cgt	gtc	gat	cag	950
Gly	Thr	Asp	Gly	Phe	Trp	Tyr	Asp	Ser	Asp	Gly	Asn	Arg	Val	Asp	Gln	
				80					85					90		
aag	acc	aat	cag	att	ctg	ctt	act	gcg	gaa	caa	ctt	aaa	aaa	aat	aac	998
Lys	Thr	Asn	Gln	Ile	Leu	Leu	Thr	Ala	Glu	Gln	Leu	Lys	Lys	Asn	Asn	
			95					100					105			
gaa	aaa	aat	tta	tca	gta	atc	agt	gat	gat	aca	tca	aaa	aaa	gat	gat	1046
Glu	Lys	Asn	Leu	Ser	Val	Ile	Ser	Asp	Asp	Thr	Ser	Lys	Lys	Asp	Asp	
		110					115					120				
gaa	aat	att	tct	aag	cag	acc	aaa	att	gct	aat	caa	caa	aca	gta	gat	1094
Glu	Asn	Ile	Ser	Lys	Gln	Thr	Lys	Ile	Ala	Asn	Gln	Gln	Thr	Val	Asp	
	125					130					135					
act	gct	aaa	ggc	ctg	act	acc	agt	aat	tta	tct	gat	ccc	atc	act	ggg	1142
Thr	Ala	Lys	Gly	Leu	Thr	Thr	Ser	Asn	Leu	Ser	Asp	Pro	Ile	Thr	Gly	
140					145					150					155	
ggt	cac	tat	gaa	aat	cac	aat	ggc	tac	ttt	gtt	tat	ata	gat	gct	tca	1190
Gly	His	Tyr	Glu	Asn	His	Asn	Gly	Tyr	Phe	Val	Tyr	Ile	Asp	Ala	Ser	
				160					165					170		
gga	aaa	caa	gta	aca	ggt	ttg	caa	aat	att	gat	ggt	aat	tta	caa	tat	1238
Gly	Lys	Gln	Val	Thr	Gly	Leu	Gln	Asn	Ile	Asp	Gly	Asn	Leu	Gln	Tyr	
			175					180					185			
ttt	gat	gac	aat	gga	tat	caa	gtc	aag	gga	tcc	ttc	cga	gat	gtc	aac	1286
Phe	Asp	Asp	Asn	Gly	Tyr	Gln	Val	Lys	Gly	Ser	Phe	Arg	Asp	Val	Asn	
		190					195					200				
ggc	aag	cat	atc	tat	ttt	gat	tca	gta	aca	ggg	aaa	gct	agt	tca	aat	1334
Gly	Lys	His	Ile	Tyr	Phe	Asp	Ser	Val	Thr	Gly	Lys	Ala	Ser	Ser	Asn	
	205					210					215					
gtt	gat	att	gtt	aac	ggt	aaa	gct	caa	gga	tat	gat	gcg	caa	ggc	aac	1382
Val	Asp	Ile	Val	Asn	Gly	Lys	Ala	Gln	Gly	Tyr	Asp	Ala	Gln	Gly	Asn	
220					225					230					235	
caa	tta	aag	aaa	agt	tat	gtc	gcc	gat	agt	tct	ggg	caa	act	tac	tat	1430
Gln	Leu	Lys	Lys	Ser	Tyr	Val	Ala	Asp	Ser	Ser	Gly	Gln	Thr	Tyr	Tyr	
				240					245					250		
ttt	gat	ggt	aat	ggc	caa	ccg	tta	atc	ggc	ttg	caa	aca	att	gat	ggg	1478
Phe	Asp	Gly	Asn	Gly	Gln	Pro	Leu	Ile	Gly	Leu	Gln	Thr	Ile	Asp	Gly	
			255					260					265			
aac	cta	caa	tat	ttt	aac	caa	caa	ggg	gtt	caa	ata	aag	ggt	ggt	ttc	1526
Asn	Leu	Gln	Tyr	Phe	Asn	Gln	Gln	Gly	Val	Gln	Ile	Lys	Gly	Gly	Phe	
		270					275					280				
caa	gat	gtt	aac	aat	aaa	cgt	att	tat	ttt	gca	cca	aac	aca	ggt	aat	1574

Gln 285	Asp	Val	Asn	Asn	Lys	Arg 290	Ile	Tyr	Phe	Ala	Pro 295	Asn	Thr	Gly	Asn	
gcc	gtt	gcc	aat	act	gaa	ata	att	aac	ggt	aaa	tta	cag	ggg	cgt	gac	1622
Ala	Val	Ala	Asn	Thr	Glu	Ile	Ile	Asn	Gly	Lys	Leu	Gln	Gly	Arg	Asp	
300					305					310					315	
gca	aat	ggt	aac	cag	gta	aag	aat	gca	ttt	agt	aaa	gat	gtt	gca	gga	1670
Ala	Asn	Gly	Asn	Gln	Val	Lys	Asn	Ala	Phe	Ser	Lys	Asp	Val	Ala	Gly	
				320					325					330		
aat	aca	ttt	tat	ttt	gac	gca	aac	ggt	gtg	atg	tta	aca	ggg	ttg	caa	1718
Asn	Thr	Phe	Tyr	Phe	Asp	Ala	Asn	Gly	Val	Met	Leu	Thr	Gly	Leu	Gln	
			335					340					345			
act	att	tca	gga	aag	aca	tat	tat	ctt	gat	gaa	caa	gga	cac	ctg	aga	1766
Thr	Ile	Ser	Gly	Lys	Thr	Tyr	Tyr	Leu	Asp	Glu	Gln	Gly	His	Leu	Arg	
		350					355					360				
aaa	aat	tac	gcg	gga	aca	ttc	aat	aat	cag	ttt	atg	tac	ttc	gat	gct	1814
Lys	Asn	Tyr	Ala	Gly	Thr	Phe	Asn	Asn	Gln	Phe	Met	Tyr	Phe	Asp	Ala	
	365					370					375					
gat	aca	ggt	gcg	ggt	aaa	aca	gcg	att	gaa	tat	caa	ttt	gat	caa	gga	1862
Asp	Thr	Gly	Ala	Gly	Lys	Thr	Ala	Ile	Glu	Tyr	Gln	Phe	Asp	Gln	Gly	
380					385				390						395	
ttg	gta	tca	caa	agt	aat	gaa	aat	act	cct	cac	aat	gcc	gca	aag	tct	1910
Leu	Val	Ser	Gln	Ser	Asn	Glu	Asn	Thr	Pro	His	Asn	Ala	Ala	Lys	Ser	
				400					405					410		
tat	gat	aaa	agt	agt	ttt	gaa	aat	gtt	gat	ggt	tac	tta	aca	gca	gat	1958
Tyr	Asp	Lys	Ser	Ser	Phe	Glu	Asn	Val	Asp	Gly	Tyr	Leu	Thr	Ala	Asp	
			415					420					425			
aca	tgg	tat	cgt	cca	acc	gat	att	tta	aaa	aat	gga	gat	act	tgg	acg	2006
Thr	Trp	Tyr	Arg	Pro	Thr	Asp	Ile	Leu	Lys	Asn	Gly	Asp	Thr	Trp	Thr	
		430					435					440				
gca	tct	acc	gaa	act	gat	atg	cgt	ccg	ctt	tta	atg	aca	tgg	tgg	cct	2054
Ala	Ser	Thr	Glu	Thr	Asp	Met	Arg	Pro	Leu	Leu	Met	Thr	Trp	Trp	Pro	
	445					450					455					
gac	aaa	caa	aca	caa	gca	aat	tac	ttg	aat	ttt	atg	tct	agt	aaa	gga	2102
Asp	Lys	Gln	Thr	Gln	Ala	Asn	Tyr	Leu	Asn	Phe	Met	Ser	Ser	Lys	Gly	
460					465					470					475	
ctt	ggt	ata	acg	acc	act	tat	aca	gca	gct	acg	tca	caa	aaa	aca	cta	2150
Leu	Gly	Ile	Thr	Thr	Thr	Tyr	Thr	Ala	Ala	Thr	Ser	Gln	Lys	Thr	Leu	
				480					485					490		
aat	gac	gca	gcc	ttt	gtt	att	caa	aca	gca	att	gaa	caa	caa	ata	tct	2198
Asn	Asp	Ala	Ala	Phe	Val	Ile	Gln	Thr	Ala	Ile	Glu	Gln	Gln	Ile	Ser	
			495					500					505			
ttg	aaa	aaa	agt	act	gag	tgg	tta	cgt	gat	gca	att	gat	agt	ttt	gtg	2246
Leu	Lys	Lys	Ser	Thr	Glu	Trp	Leu	Arg	Asp	Ala	Ile	Asp	Ser	Phe	Val	
		510					515					520				
aag	acg	caa	gct	aat	tgg	aat	aag	caa	aca	gaa	gat	gaa	gct	ttc	gat	2294

Lys 525	Thr	Gln	Ala	Asn	Trp	Asn 530	Lys	Gln	Thr	Glu	Asp 535	Glu	Ala	Phe	Asp	
ggt	ttg	cag	tggt	ctt	caa	ggg	gga	ttc	cta	gct	tat	caa	gat	gat	tca	2342
Gly 540	Leu	Gln	Trp	Leu	Gln	Gly	Gly	Phe	Leu	Ala	Tyr	Gln	Asp	Asp	Ser 555	
cat	cgg	acg	ccg	aat	act	gat	tca	gga	aat	aac	aga	aaa	cta	gga	cgt	2390
His	Arg	Thr	Pro	Asn 560	Thr	Asp	Ser	Gly	Asn 565	Asn	Arg	Lys	Leu	Gly 570	Arg	
caa	cca	att	aat	atc	gat	ggt	tcg	aaa	gat	aca	act	gat	ggt	aaa	ggc	2438
Gln	Pro	Ile	Asn 575	Ile	Asp	Gly	Ser	Lys	Asp 580	Thr	Thr	Asp	Gly 585	Lys	Gly	
tct	gaa	ttc	tta	tta	gct	aac	gat	att	gac	aac	tca	aat	ccg	att	gtt	2486
Ser	Glu	Phe 590	Leu	Leu	Ala	Asn	Asp 595	Ile	Asp	Asn	Ser	Asn 600	Pro	Ile	Val	
caa	gct	gag	caa	tta	aac	tggt	cta	cac	tat	tta	atg	aat	ttt	ggt	agt	2534
Gln	Ala 605	Glu	Gln	Leu	Asn	Trp 610	Leu	His	Tyr	Leu	Met 615	Asn	Phe	Gly	Ser	
att	aca	ggt	aat	aat	gac	aat	gcg	aat	ttt	gat	ggc	att	cgt	gta	gat	2582
Ile 620	Thr	Gly	Asn	Asn	Asp 625	Asn	Ala	Asn	Phe	Asp 630	Gly	Ile	Arg	Val	Asp 635	
gct	gtt	gat	aat	gtt	gat	gct	gat	tta	cta	aaa	ata	gct	ggc	gat	tat	2630
Ala	Val	Asp	Asn	Val 640	Asp	Ala	Asp	Leu	Leu 645	Lys	Ile	Ala	Gly	Asp 650	Tyr	
ttt	aaa	gct	cta	tat	ggt	aca	gat	aaa	agc	gac	gcc	aat	gcc	aat	aag	2678
Phe	Lys	Ala	Leu 655	Tyr	Gly	Thr	Asp 660	Lys	Ser	Asp	Ala	Asn 665	Ala	Asn	Lys	
cat	ttg	tct	att	tta	gaa	gac	tggt	aac	ggt	aaa	gat	cct	cag	tat	gtt	2726
His	Leu 670	Ser	Ile	Leu	Glu	Asp	Trp 675	Asn	Gly	Lys	Asp 680	Pro	Gln	Tyr	Val	
aat	caa	cag	ggc	aat	gcg	caa	tta	aca	atg	gat	tac	aca	gtt	act	tca	2774
Asn	Gln 685	Gln	Gly	Asn	Ala	Gln 690	Leu	Thr	Met	Asp 695	Tyr	Thr	Val	Thr	Ser	
cag	ttt	ggc	aat	tct	cta	aca	cat	ggc	gcc	aac	aac	agg	agt	aac	atg	2822
Gln 700	Phe	Gly	Asn	Ser	Leu 705	Thr	His	Gly	Ala	Asn 710	Asn	Arg	Ser	Asn	Met 715	
tggt	tat	ttc	tta	gat	act	ggc	tat	tat	ctt	aat	gga	gat	ctt	aat	aag	2870
Trp	Tyr	Phe	Leu	Asp 720	Thr	Gly	Tyr	Tyr	Leu 725	Asn	Gly	Asp	Leu	Asn 730	Lys	
aag	ata	gta	gat	aag	aac	cgt	cca	aat	tct	ggc	act	ttg	gtt	aac	aga	2918
Lys	Ile	Val	Asp 735	Lys	Asn	Arg	Pro	Asn 740	Ser	Gly	Thr	Leu	Val	Asn 745	Arg	
att	gct	aat	tca	ggt	gat	aca	aaa	gtt	att	cca	aat	tat	agt	ttt	gtt	2966
Ile	Ala	Asn	Ser 750	Gly	Asp	Thr	Lys 755	Val	Ile	Pro	Asn 760	Tyr	Ser	Phe	Val	
aga	gca	cat	gat	tac	gat	gct	caa	gat	cca	att	aga	aaa	gcc	atg	att	3014

Arg 765	Ala	His	Asp	Tyr	Asp	Ala 770	Gln	Asp	Pro	Ile	Arg 775	Lys	Ala	Met	Ile	
gat	cat	ggt	att	att	aaa	aac	atg	cag	gat	act	ttc	act	ttt	gac	caa	3062
Asp 780	His	Gly	Ile	Ile	Lys 785	Asn	Met	Gln	Asp	Thr 790	Phe	Thr	Phe	Asp	Gln 795	
ctg	gct	cag	gga	atg	gaa	ttc	tac	tat	aaa	gat	caa	gag	aat	ccg	tct	3110
Leu	Ala	Gln	Gly	Met 800	Glu	Phe	Tyr	Tyr	Lys 805	Asp	Gln	Glu	Asn	Pro 810	Ser	
ggt	ttc	aaa	aag	tat	aac	gat	tat	aac	tta	cct	agt	gct	tat	gca	atg	3158
Gly	Phe	Lys	Lys 815	Tyr	Asn	Asp	Tyr	Asn 820	Leu	Pro	Ser	Ala	Tyr 825	Ala	Met	
ttg	ttg	act	aat	aag	gat	act	gta	cct	cgt	gtc	tat	tat	gga	gat	atg	3206
Leu	Leu	Thr 830	Asn	Lys	Asp	Thr	Val 835	Pro	Arg	Val	Tyr 840	Tyr	Gly	Asp	Met	
tac	ctc	gaa	ggc	ggg	caa	tat	atg	gaa	aaa	ggg	acg	att	tac	aat	cct	3254
Tyr	Leu	Glu	Gly	Gly	Gln 850	Tyr	Met	Glu	Lys	Gly	Thr 855	Ile	Tyr	Asn	Pro	
gtc	att	tca	gcg	ttg	ctc	aaa	gct	aga	ata	aaa	tat	ggt	tct	ggt	ggg	3302
Val 860	Ile	Ser	Ala	Leu	Leu 865	Lys	Ala	Arg	Ile	Lys 870	Tyr	Val	Ser	Gly	Gly 875	
caa	aca	atg	gct	acc	gat	agt	tct	gga	aaa	gac	ctt	aaa	gat	ggc	gaa	3350
Gln	Thr	Met	Ala	Thr 880	Asp	Ser	Ser	Gly	Lys 885	Asp	Leu	Lys	Asp	Gly 890	Glu	
act	gat	ttg	tta	aca	agt	ggt	cga	ttt	ggt	aaa	gga	att	atg	aca	tca	3398
Thr	Asp	Leu	Leu 895	Thr	Ser	Val	Arg	Phe 900	Gly	Lys	Gly	Ile	Met 905	Thr	Ser	
gat	caa	acc	aca	aca	caa	gac	aat	agc	caa	gat	tat	aaa	aat	caa	ggc	3446
Asp	Gln	Thr 910	Thr	Thr	Gln	Asp	Asn 915	Ser	Gln	Asp	Tyr	Lys 920	Asn	Gln	Gly	
atc	ggt	gtc	att	ggt	ggt	aat	aac	cct	gac	ctt	aag	ttg	aac	aat	gat	3494
Ile	Gly 925	Val	Ile	Val	Gly	Asn 930	Asn	Pro	Asp	Leu	Lys 935	Leu	Asn	Asn	Asp	
aag	acc	att	acc	ttg	cat	atg	gga	aag	gcg	cat	aag	aat	caa	ctt	tac	3542
Lys 940	Thr	Ile	Thr	Leu	His 945	Met	Gly	Lys	Ala	His 950	Lys	Asn	Gln	Leu	Tyr 955	
cgt	gcc	tta	gta	tta	tca	aat	gac	tca	gga	att	gat	ggt	tat	gat	agt	3590
Arg	Ala	Leu	Val	Leu 960	Ser	Asn	Asp	Ser	Gly	Ile	Asp	Val	Tyr	Asp 970	Ser	
gat	gat	aaa	gca	cca	act	ttg	aga	aca	aat	gac	aac	ggt	gac	ttg	att	3638
Asp	Asp	Lys	Ala 975	Pro	Thr	Leu	Arg	Thr 980	Asn	Asp	Asn	Gly	Asp 985	Leu	Ile	
ttc	cat	aag	aca	aat	acg	ttt	gtg	aag	caa	gat	gga	act	att	ata	aat	3686
Phe	His	Lys 990	Thr	Asn	Thr	Phe	Val 995	Lys	Gln	Asp	Gly	Thr 1000	Ile	Ile	Asn	
tac	gaa	atg	aag	gga	tca	tta	aat	gct	tta	att	tca	ggt	tat	tta	ggt	3734

Tyr Glu Met Lys Gly Ser Leu Asn Ala Leu Ile Ser Gly Tyr Leu Gly	
1005 1010 1015	
gtc tgg gtg cca gtt gga gct agt gat tca caa gat gct cgt aca gtg	3782
Val Trp Val Pro Val Gly Ala Ser Asp Ser Gln Asp Ala Arg Thr Val	
1020 1025 1030 1035	
gca act gag tca tca tca agt aat gat ggt tct gta ttc cat tca aat	3830
Ala Thr Glu Ser Ser Ser Ser Asn Asp Gly Ser Val Phe His Ser Asn	
1040 1045 1050	
gct gca tta gat tct aat gtt ata tat gaa ggc ttt tca aac ttt caa	3878
Ala Ala Leu Asp Ser Asn Val Ile Tyr Glu Gly Phe Ser Asn Phe Gln	
1055 1060 1065	
gcg atg ccg act tct cct gag caa agt aca aat gtt gtt att gca aca	3926
Ala Met Pro Thr Ser Pro Glu Gln Ser Thr Asn Val Val Ile Ala Thr	
1070 1075 1080	
aag gct aac tta ttt aaa gaa tta ggt att act agt ttt gag tta gca	3974
Lys Ala Asn Leu Phe Lys Glu Leu Gly Ile Thr Ser Phe Glu Leu Ala	
1085 1090 1095	
cct caa tat agg tct agt ggt gac act aat tac ggt ggc atg tca ttc	4022
Pro Gln Tyr Arg Ser Ser Gly Asp Thr Asn Tyr Gly Gly Met Ser Phe	
1100 1105 1110 1115	
tta gat tct ttc tta aat aat ggt tat gca ttt acc gat aga tat gat	4070
Leu Asp Ser Phe Leu Asn Asn Gly Tyr Ala Phe Thr Asp Arg Tyr Asp	
1120 1125 1130	
tta ggc ttt aac aaa gca gac ggg aat cct aac cca aca aag tat gga	4118
Leu Gly Phe Asn Lys Ala Asp Gly Asn Pro Asn Pro Thr Lys Tyr Gly	
1135 1140 1145	
aca gat caa gat tta cgt aat gca ata gag gca tta cac aaa aac ggc	4166
Thr Asp Gln Asp Leu Arg Asn Ala Ile Glu Ala Leu His Lys Asn Gly	
1150 1155 1160	
atg cag gct ata gct gat tgg gtt cct gac caa ata tat gct tta cca	4214
Met Gln Ala Ile Ala Asp Trp Val Pro Asp Gln Ile Tyr Ala Leu Pro	
1165 1170 1175	
gga aag gaa gtt gtt acc gct act aga gta gac gaa cgg gga aat caa	4262
Gly Lys Glu Val Val Thr Ala Thr Arg Val Asp Glu Arg Gly Asn Gln	
1180 1185 1190 1195	
cta aaa gac aca gat ttt gtc aac tta ctc tat gtt gct aat act aaa	4310
Leu Lys Asp Thr Asp Phe Val Asn Leu Leu Tyr Val Ala Asn Thr Lys	
1200 1205 1210	
agt agt ggt gtg gat tat cag gca aag tat ggc ggc gaa ttt tta gat	4358
Ser Ser Gly Val Asp Tyr Gln Ala Lys Tyr Gly Gly Glu Phe Leu Asp	
1215 1220 1225	
aaa tta aga gaa gag tac cca tcg tta ttc aaa cag aac caa gta tcg	4406
Lys Leu Arg Glu Glu Tyr Pro Ser Leu Phe Lys Gln Asn Gln Val Ser	
1230 1235 1240	
aca ggt cag cca att gat gct tct aca aaa att aag caa tgg tca gct	4454

Thr Gly Gln Pro Ile Asp Ala Ser Thr Lys Ile Lys Gln Trp Ser Ala	
1245 1250 1255	
aaa tat atg aat ggg acc aat att tta cat cga ggt gct tat tat gtt	4502
Lys Tyr Met Asn Gly Thr Asn Ile Leu His Arg Gly Ala Tyr Tyr Val	
1260 1265 1270 1275	
ttg aaa gac tgg gct act aac cag tat ttt aac att gca aaa acg aat	4550
Leu Lys Asp Trp Ala Thr Asn Gln Tyr Phe Asn Ile Ala Lys Thr Asn	
1280 1285 1290	
gaa gta ttt ttg cca cta cag ttg cag aat aaa gat gcg caa act ggt	4598
Glu Val Phe Leu Pro Leu Gln Leu Gln Asn Lys Asp Ala Gln Thr Gly	
1295 1300 1305	
ttc att agt gat gcc tcc ggt gta aaa tat tac tca att agt ggt tat	4646
Phe Ile Ser Asp Ala Ser Gly Val Lys Tyr Tyr Ser Ile Ser Gly Tyr	
1310 1315 1320	
caa gca aaa gat act ttt att gaa gat ggt aat ggg aat tgg tat tac	4694
Gln Ala Lys Asp Thr Phe Ile Glu Asp Gly Asn Gly Asn Trp Tyr Tyr	
1325 1330 1335	
ttt gat aaa gat ggt tac atg gtg cgt tcg cag caa gga gaa aat cct	4742
Phe Asp Lys Asp Gly Tyr Met Val Arg Ser Gln Gln Gly Glu Asn Pro	
1340 1345 1350 1355	
ata aga aca gtc gaa act agt gtc aac aca cga aac ggt aat tat tac	4790
Ile Arg Thr Val Glu Thr Ser Val Asn Thr Arg Asn Gly Asn Tyr Tyr	
1360 1365 1370	
ttt atg cca aat ggt gtc gag ttg cgc aaa ggc ttt gga acg gat aat	4838
Phe Met Pro Asn Gly Val Glu Leu Arg Lys Gly Phe Gly Thr Asp Asn	
1375 1380 1385	
agt ggt aat gtc tat tat ttt gat gat caa ggt aag atg gtg aga gat	4886
Ser Gly Asn Val Tyr Tyr Phe Asp Asp Gln Gly Lys Met Val Arg Asp	
1390 1395 1400	
aaa tac att aac gat gat gct aat aat ttt tat cac tta aat gtt gat	4934
Lys Tyr Ile Asn Asp Asp Ala Asn Asn Phe Tyr His Leu Asn Val Asp	
1405 1410 1415	
ggg act atg tct cga gga cta ttt aaa ttt gat tct gat act cta cag	4982
Gly Thr Met Ser Arg Gly Leu Phe Lys Phe Asp Ser Asp Thr Leu Gln	
1420 1425 1430 1435	
tat ttt gct agt aat ggt gtc caa ata aaa gat agt tat gcg aag gat	5030
Tyr Phe Ala Ser Asn Gly Val Gln Ile Lys Asp Ser Tyr Ala Lys Asp	
1440 1445 1450	
agt aaa ggc aat aaa tat tat ttt gac tca gct aca gga aat aac gat	5078
Ser Lys Gly Asn Lys Tyr Tyr Phe Asp Ser Ala Thr Gly Asn Asn Asp	
1455 1460 1465	
act ggg aaa gcc caa act tgg gat ggt aat ggc tac tat att act att	5126
Thr Gly Lys Ala Gln Thr Trp Asp Gly Asn Gly Tyr Tyr Ile Thr Ile	
1470 1475 1480	
gat tct gat gcg aac aat aca att ggg gtt aac aca gac tac act gcc	5174

Asp Ser Asp Ala Asn Asn Thr Ile Gly Val Asn Thr Asp Tyr Thr Ala	
1485 1490 1495	
tac atc act agc tcg ctg cgc gaa gat ggc tta ttt gct aac gca cct	5222
Tyr Ile Thr Ser Ser Leu Arg Glu Asp Gly Leu Phe Ala Asn Ala Pro	
1500 1505 1510 1515	
tac ggt gtt gta aca aaa gac caa aat ggt aac gat ctt aag tgg cag	5270
Tyr Gly Val Val Thr Lys Asp Gln Asn Gly Asn Asp Leu Lys Trp Gln	
1520 1525 1530	
tat att aac cat acg aaa cag tac gaa ggg caa caa gtg caa gtc acg	5318
Tyr Ile Asn His Thr Lys Gln Tyr Glu Gly Gln Gln Val Gln Val Thr	
1535 1540 1545	
cgt caa tac aca gac agt aag gga gtc agc tgg aac tta att acc ttt	5366
Arg Gln Tyr Thr Asp Ser Lys Gly Val Ser Trp Asn Leu Ile Thr Phe	
1550 1555 1560	
gct ggt ggt gat tta caa gga caa agg ctt tgg gtg gat agt cgt gcg	5414
Ala Gly Gly Asp Leu Gln Gly Gln Arg Leu Trp Val Asp Ser Arg Ala	
1565 1570 1575	
tta act atg aca cca ttt aaa acg atg aac caa ata agc ttc att agt	5462
Leu Thr Met Thr Pro Phe Lys Thr Met Asn Gln Ile Ser Phe Ile Ser	
1580 1585 1590 1595	
tat gct aac cgc aat gat ggg ttg ttt ttg aat gcg cca tac caa gtc	5510
Tyr Ala Asn Arg Asn Asp Gly Leu Phe Leu Asn Ala Pro Tyr Gln Val	
1600 1605 1610	
aag ggg tat caa tta gct ggg atg tcc aac caa tac aag ggc caa caa	5558
Lys Gly Tyr Gln Leu Ala Gly Met Ser Asn Gln Tyr Lys Gly Gln Gln	
1615 1620 1625	
gtg acc att gct ggg gtg gcg aac gtt tct gga aaa gac tgg agt ctg	5606
Val Thr Ile Ala Gly Val Ala Asn Val Ser Gly Lys Asp Trp Ser Leu	
1630 1635 1640	
att agt ttt aat ggg aca cag tac tgg att gat agt cag gca ttg aat	5654
Ile Ser Phe Asn Gly Thr Gln Tyr Trp Ile Asp Ser Gln Ala Leu Asn	
1645 1650 1655	
acc aat ttc aca cat gac atg aac caa aag gtc ttt gtc aat aca act	5702
Thr Asn Phe Thr His Asp Met Asn Gln Lys Val Phe Val Asn Thr Thr	
1660 1665 1670 1675	
agt aat ctt gat ggg tta ttc tta aat gcg cca tac cgt caa ccg ggt	5750
Ser Asn Leu Asp Gly Leu Phe Leu Asn Ala Pro Tyr Arg Gln Pro Gly	
1680 1685 1690	
tat aag tta gcc ggt ttg gct aaa aat tac aac aac caa acg gtt act	5798
Tyr Lys Leu Ala Gly Leu Ala Lys Asn Tyr Asn Asn Gln Thr Val Thr	
1695 1700 1705	
gtt agt caa cag tac ttt gat gat caa ggc acg gtc tgg agt cag gtt	5846
Val Ser Gln Gln Tyr Phe Asp Asp Gln Gly Thr Val Trp Ser Gln Val	
1710 1715 1720	
gtc ctt ggg ggt cag acg gtc tgg gtt gat aac cat gca ttg gca cag	5894

9/36

Val 1725	Leu	Gly	Gly	Gln	Thr	Val	Trp	Val	Asp	Asn	His	Ala	Leu	Ala	Gln	
1730																
atg	caa	gtt	agt	gat	aca	gac	caa	cag	ctc	tat	gtg	aat	agc	aat	ggt	5942
Met	Gln	Val	Ser	Asp	Thr	Asp	Gln	Gln	Leu	Tyr	Val	Asn	Ser	Asn	Gly	
1740																
1745																
1750																
cgg	aat	gat	ggg	tta	ttc	ttg	aat	gcg	cca	tat	cgt	ggt	caa	ggg	tca	5990
Arg	Asn	Asp	Gly	Leu	Phe	Leu	Asn	Ala	Pro	Tyr	Arg	Gly	Gln	Gly	Ser	
1760																
1765																
1770																
caa	ctg	ata	ggc	atg	acg	gca	gat	tat	aat	ggg	caa	cat	gta	caa	gtg	6038
Gln	Leu	Ile	Gly	Met	Thr	Ala	Asp	Tyr	Asn	Gly	Gln	His	Val	Gln	Val	
1775																
1780																
1785																
acc	aag	caa	ggg	caa	gat	gcc	tat	ggt	gca	caa	tgg	cgt	ctt	att	acg	6086
Thr	Lys	Gln	Gly	Gln	Asp	Ala	Tyr	Gly	Ala	Gln	Trp	Arg	Leu	Ile	Thr	
1790																
1795																
1800																
cta	aat	aat	caa	cag	gtc	tgg	gtt	gat	agt	cgc	gct	ttg	agc	aca	aca	6134
Leu	Asn	Asn	Gln	Gln	Val	Trp	Val	Asp	Ser	Arg	Ala	Leu	Ser	Thr	Thr	
1805																
1810																
1815																
atc	atg	caa	gcc	atg	aat	gat	aat	atg	tat	gta	aat	agc	agc	caa	cgg	6182
Ile	Met	Gln	Ala	Met	Asn	Asp	Asn	Met	Tyr	Val	Asn	Ser	Ser	Gln	Arg	
1820																
1825																
1830																
1835																
aca	gat	ggc	ttg	tgg	tta	aac	gca	cct	tat	acg	atg	agt	ggg	gct	aaa	6230
Thr	Asp	Gly	Leu	Trp	Leu	Asn	Ala	Pro	Tyr	Thr	Met	Ser	Gly	Ala	Lys	
1840																
1845																
1850																
tgg	gct	ggt	gat	aca	cgt	tca	gct	aat	ggg	cgc	tat	gtc	cat	att	tca	6278
Trp	Ala	Gly	Asp	Thr	Arg	Ser	Ala	Asn	Gly	Arg	Tyr	Val	His	Ile	Ser	
1855																
1860																
1865																
aaa	gct	tat	tca	aac	gaa	gtc	ggc	aat	aca	tat	tac	ttg	acg	aat	ttg	6326
Lys	Ala	Tyr	Ser	Asn	Glu	Val	Gly	Asn	Thr	Tyr	Tyr	Leu	Thr	Asn	Leu	
1870																
1875																
1880																
aat	ggt	caa	agc	aca	tgg	att	gac	aag	cgg	gcg	ttt	act	gtg	acc	ttc	6374
Asn	Gly	Gln	Ser	Thr	Trp	Ile	Asp	Lys	Arg	Ala	Phe	Thr	Val	Thr	Phe	
1885																
1890																
1895																
gat	cag	gtg	gtg	gca	tta	aat	gca	acg	att	gtg	gca	cgc	caa	cga	cca	6422
Asp	Gln	Val	Val	Ala	Leu	Asn	Ala	Thr	Ile	Val	Ala	Arg	Gln	Arg	Pro	
1900																
1905																
1910																
1915																
gat	ggg	atg	ttt	aag	aca	gca	cca	tat	ggt	gaa	gcg	ggg	gcg	cag	ttt	6470
Asp	Gly	Met	Phe	Lys	Thr	Ala	Pro	Tyr	Gly	Glu	Ala	Gly	Ala	Gln	Phe	
1920																
1925																
1930																
gtc	gat	tat	gtg	aca	aac	tat	aac	cag	caa	acc	gtg	cca	gta	aca	aag	6518
Val	Asp	Tyr	Val	Thr	Asn	Tyr	Asn	Gln	Gln	Thr	Val	Pro	Val	Thr	Lys	
1935																
1940																
1945																
caa	cat	tca	gat	gct	cag	ggg	aat	caa	tgg	tac	tta	gcg	aca	gtg	aat	6566
Gln	His	Ser	Asp	Ala	Gln	Gly	Asn	Gln								

Gly Thr Gln Tyr Trp Ile Asp Gln Arg Ser Phe Ser Pro Val Val Thr	
1965 1970 1975	
aag gtg gtt gat tat caa gct aag att gtg cca cgg aca aca cgt gat	6662
Lys Val Val Asp Tyr Gln Ala Lys Ile Val Pro Arg Thr Thr Arg Asp	
1980 1985 1990 1995	
ggt gtg ttt agt ggc gca ccc tat ggg gaa gtg aat gct aag cta gtt	6710
Gly Val Phe Ser Gly Ala Pro Tyr Gly Glu Val Asn Ala Lys Leu Val	
2000 2005 2010	
aac atg gca act gcg tat caa aat caa gtt gtc cat gcg aca ggg gaa	6758
Asn Met Ala Thr Ala Tyr Gln Asn Gln Val Val His Ala Thr Gly Glu	
2015 2020 2025	
tat acg aat gct tca ggg atc aca tgg agt cag ttc gcg tta agc ggg	6806
Tyr Thr Asn Ala Ser Gly Ile Thr Trp Ser Gln Phe Ala Leu Ser Gly	
2030 2035 2040	
caa gaa gac aag cta tgg att gat aag cgt gct ttg caa gct	6848
Gln Glu Asp Lys Leu Trp Ile Asp Lys Arg Ala Leu Gln Ala	
2045 2050 2055	
taaggggaagg attcgacaaa ggaggggtaac attatcagcg gatggtgtta tctctcctttc	6908
ctgtactcag tatttcccaa ataattgaga cagtttcatg acaaatcaac aaaactagtg	6968
tcaatgcctc ggttatgggg taaactacta ttagttaaag ggttggtgca taataatatac	7028
acattcaata tattatgtat ttttatctga ttatgtgatt ttttgagatt tggagcgaaa	7088
atgaataaag taataattaa taaaagtttc tgtgttttgg taatgagttt gttgtcgata	7148
tttctattct cattaagagt agacgctcgt tccaatagtg gctacaatca aatatctttt	7208
aatcaacatg aaattgctta tgcgcctttt agtcaagtac cttggtatgc aactgtgaat	7268
gtcggaatgt ctgcagacaa aaataatatc tatacagcta ttgatatggg gccatagtat	7328
caaggaactt ctttttttcc gtgggtggtat caacatgata attatgatta tcatgatggg	7388
tatcaacttg cgattgacaa tcaatcgctt ttcatagaaa tggttgatac aggaaggtcg	7448
tcgtttgatg ttattgagtt gtatacaggt aaaaaagttg gttatggaaa attcacagcc	7508
tataactata acaagacttc atggcagttt caaatatctt taaaggcgat aaagaaaatt	7568
ttaaataata acatatcaga aaatgcgaca attagtttgt acaatggtta tctattcaaa	7628
caagccgaaa gtatcacgta tgctgggttca ccaacagggc catggttact tgccggtatt	7688
ggtttggttt ttgctgggtg gggctattac ctgcaaaatc agcgccgaaa aacgattacg	7748
gcatcatcaa caaacgaggt aacctatgct taataatgct gttattattg gtctaattat	7808
tatttggttc tatgtattat ctgttctgaa gcgtgctaaa acggatgcgt tttatttttt	7868
ggctggcagc gcaggcttgt tcgtcattct tcttttatgg agtaaacctt atgggtgttg	7928
gttgttttcg accattctca cttggtcggc cggggttggt ggtcatctca ctggcttatt	7988

tgatacttttc tatgcgtcac atgttattca ggttgtggcg aatcatcata ccagcatttt 8048
 atttggttgat tatgaatggt caggcattat tgaaacaacg gcattttggg gattaattgc 8108
 cttttatcct gtgtatgatg cacaaaagcg gttaatgttg gcgcttattg gggcattatg 8168
 gtttttcctt gcaaatacgt tacgactatc ttttgttgca gttgtaattt attatttttg 8228
 cgatggcgct ttttatgttg ctcatccaat tatcggacgc ctgttgttct attcaattgt 8288
 aatcttgctt tattatctag tcttcataaa ggggcaactt gttaatcaaa tgctcaaaaa 8348
 acagggataa actatgtcat tttatttaca tcaatttact tatcagttag gtttttggtt 8408
 cacttggttg ctcatccgt ttgttgtgga aattttccca gctattattt caatctagtt 8468
 ttgataaaaa catcaaaaaa gcatcaagtg atgcaagaac cattgaaatt accaatggta 8528
 tctattgtat tacctatata taattcagga caaacattat accagtgtat tcagtcgatt 8588
 agtcaatcaa cctatccaaa acaattaatc caaattattg ccgtgaataa tcaaagcact 8648
 gataacagtt ttacagtgtt taatcaggct caagctgatt tccaataact tagaatgcag 8708
 tggatgaaca cagatcaagg taaagcgcggt gcactaaatg ctgctattta taatagtatg 8768
 gggcaatata ttattaattt ggatactgat ggttggttag aaccatgc cttaaaacgg 8828
 tttgtgcttt atttgaaaa tcaactcagaa attgatgtag caactggtac gattctgaca 8888
 caaaaaaaaa tgattcaaaa aacgcaaagt aaatggctta aattgctaca gttaaacgaa 8948
 tattttgagt atgcacaatc attcttatca ggccgtagta ttgaaaaccg tggtaatcgt 9008
 ttgttcacaa tgtcagggtc attttcagcg ttttagacgtg atgtattagt tcagacattt 9068
 atgtataatg ttgacacggt tggatgaagat actgatatga catttcaact gcgattccgc 9128
 ctaggcaagc gtattggttt ctgtgatgat gccatgtttt atgttgaacc aatatcaggt 9188
 tatagtgaac tttatttaca acggcagcgt tggcagcgtg ggcaaattga agtcgcacaa 9248
 aatttcatgc agaataagtt gagtgtccgt cagattttta ctaactttat gattagtcga 9308
 ttaatgattg atc 9321

<210> 2

<211> 2057

<212> PRT

<213> *Leuconostoc mesenteroides*

<400> 2

Met Lys Gln Gln Glu Thr Val Thr Arg Lys Lys Leu Tyr Lys Ser Gly
 1 5 10 15

Lys Val Trp Val Ala Ala Ala Thr Ala Phe Ala Val Leu Gly Val Ser
 20 25 30

12/36

Thr Val Thr Thr Val His Ala Asp Thr Asn Ser Asn Val Ala Val Lys
 35 40 45
 Gln Ile Asn Asn Thr Gly Thr Asn Asp Ser Gly Glu Lys Lys Val Pro
 50 55 60
 Val Pro Ser Thr Asn Asn Asp Ser Leu Lys Gln Gly Thr Asp Gly Phe
 65 70 75 80
 Trp Tyr Asp Ser Asp Gly Asn Arg Val Asp Gln Lys Thr Asn Gln Ile
 85 90 95
 Leu Leu Thr Ala Glu Gln Leu Lys Lys Asn Asn Glu Lys Asn Leu Ser
 100 105 110
 Val Ile Ser Asp Asp Thr Ser Lys Lys Asp Asp Glu Asn Ile Ser Lys
 115 120 125
 Gln Thr Lys Ile Ala Asn Gln Gln Thr Val Asp Thr Ala Lys Gly Leu
 130 135 140
 Thr Thr Ser Asn Leu Ser Asp Pro Ile Thr Gly Gly His Tyr Glu Asn
 145 150 155 160
 His Asn Gly Tyr Phe Val Tyr Ile Asp Ala Ser Gly Lys Gln Val Thr
 165 170 175
 Gly Leu Gln Asn Ile Asp Gly Asn Leu Gln Tyr Phe Asp Asp Asn Gly
 180 185 190
 Tyr Gln Val Lys Gly Ser Phe Arg Asp Val Asn Gly Lys His Ile Tyr
 195 200 205
 Phe Asp Ser Val Thr Gly Lys Ala Ser Ser Asn Val Asp Ile Val Asn
 210 215 220
 Gly Lys Ala Gln Gly Tyr Asp Ala Gln Gly Asn Gln Leu Lys Lys Ser
 225 230 235 240
 Tyr Val Ala Asp Ser Ser Gly Gln Thr Tyr Tyr Phe Asp Gly Asn Gly
 245 250 255
 Gln Pro Leu Ile Gly Leu Gln Thr Ile Asp Gly Asn Leu Gln Tyr Phe
 260 265 270
 Asn Gln Gln Gly Val Gln Ile Lys Gly Gly Phe Gln Asp Val Asn Asn
 275 280 285
 Lys Arg Ile Tyr Phe Ala Pro Asn Thr Gly Asn Ala Val Ala Asn Thr
 290 295 300
 Glu Ile Ile Asn Gly Lys Leu Gln Gly Arg Asp Ala Asn Gly Asn Gln
 305 310 315 320
 Val Lys Asn Ala Phe Ser Lys Asp Val Ala Gly Asn Thr Phe Tyr Phe
 325 330 335
 Asp Ala Asn Gly Val Met Leu Thr Gly Leu Gln Thr Ile Ser Gly Lys
 340 345 350

Thr Tyr Tyr Leu Asp Glu Gln Gly His Leu Arg Lys Asn Tyr Ala Gly
 355 360 365
 Thr Phe Asn Asn Gln Phe Met Tyr Phe Asp Ala Asp Thr Gly Ala Gly
 370 375 380
 Lys Thr Ala Ile Glu Tyr Gln Phe Asp Gln Gly Leu Val Ser Gln Ser
 385 390 395 400
 Asn Glu Asn Thr Pro His Asn Ala Ala Lys Ser Tyr Asp Lys Ser Ser
 405 410 415
 Phe Glu Asn Val Asp Gly Tyr Leu Thr Ala Asp Thr Trp Tyr Arg Pro
 420 425 430
 Thr Asp Ile Leu Lys Asn Gly Asp Thr Trp Thr Ala Ser Thr Glu Thr
 435 440 445
 Asp Met Arg Pro Leu Leu Met Thr Trp Trp Pro Asp Lys Gln Thr Gln
 450 455 460
 Ala Asn Tyr Leu Asn Phe Met Ser Ser Lys Gly Leu Gly Ile Thr Thr
 465 470 475 480
 Thr Tyr Thr Ala Ala Thr Ser Gln Lys Thr Leu Asn Asp Ala Ala Phe
 485 490 495
 Val Ile Gln Thr Ala Ile Glu Gln Gln Ile Ser Leu Lys Lys Ser Thr
 500 505 510
 Glu Trp Leu Arg Asp Ala Ile Asp Ser Phe Val Lys Thr Gln Ala Asn
 515 520 525
 Trp Asn Lys Gln Thr Glu Asp Glu Ala Phe Asp Gly Leu Gln Trp Leu
 530 535 540
 Gln Gly Gly Phe Leu Ala Tyr Gln Asp Asp Ser His Arg Thr Pro Asn
 545 550 555 560
 Thr Asp Ser Gly Asn Asn Arg Lys Leu Gly Arg Gln Pro Ile Asn Ile
 565 570 575
 Asp Gly Ser Lys Asp Thr Thr Asp Gly Lys Gly Ser Glu Phe Leu Leu
 580 585 590
 Ala Asn Asp Ile Asp Asn Ser Asn Pro Ile Val Gln Ala Glu Gln Leu
 595 600 605
 Asn Trp Leu His Tyr Leu Met Asn Phe Gly Ser Ile Thr Gly Asn Asn
 610 615 620
 Asp Asn Ala Asn Phe Asp Gly Ile Arg Val Asp Ala Val Asp Asn Val
 625 630 635 640
 Asp Ala Asp Leu Leu Lys Ile Ala Gly Asp Tyr Phe Lys Ala Leu Tyr
 645 650 655
 Gly Thr Asp Lys Ser Asp Ala Asn Ala Asn Lys His Leu Ser Ile Leu
 660 665 670

Glu Asp Trp Asn Gly Lys Asp Pro Gln Tyr Val Asn Gln Gln Gly Asn
 675 680 685
 Ala Gln Leu Thr Met Asp Tyr Thr Val Thr Ser Gln Phe Gly Asn Ser
 690 695 700
 Leu Thr His Gly Ala Asn Asn Arg Ser Asn Met Trp Tyr Phe Leu Asp
 705 710 715 720
 Thr Gly Tyr Tyr Leu Asn Gly Asp Leu Asn Lys Lys Ile Val Asp Lys
 725 730 735
 Asn Arg Pro Asn Ser Gly Thr Leu Val Asn Arg Ile Ala Asn Ser Gly
 740 745 750
 Asp Thr Lys Val Ile Pro Asn Tyr Ser Phe Val Arg Ala His Asp Tyr
 755 760 765
 Asp Ala Gln Asp Pro Ile Arg Lys Ala Met Ile Asp His Gly Ile Ile
 770 775 780
 Lys Asn Met Gln Asp Thr Phe Thr Phe Asp Gln Leu Ala Gln Gly Met
 785 790 795 800
 Glu Phe Tyr Tyr Lys Asp Gln Glu Asn Pro Ser Gly Phe Lys Lys Tyr
 805 810 815
 Asn Asp Tyr Asn Leu Pro Ser Ala Tyr Ala Met Leu Leu Thr Asn Lys
 820 825 830
 Asp Thr Val Pro Arg Val Tyr Tyr Gly Asp Met Tyr Leu Glu Gly Gly
 835 840 845
 Gln Tyr Met Glu Lys Gly Thr Ile Tyr Asn Pro Val Ile Ser Ala Leu
 850 855 860
 Leu Lys Ala Arg Ile Lys Tyr Val Ser Gly Gly Gln Thr Met Ala Thr
 865 870 875 880
 Asp Ser Ser Gly Lys Asp Leu Lys Asp Gly Glu Thr Asp Leu Leu Thr
 885 890 895
 Ser Val Arg Phe Gly Lys Gly Ile Met Thr Ser Asp Gln Thr Thr Thr
 900 905 910
 Gln Asp Asn Ser Gln Asp Tyr Lys Asn Gln Gly Ile Gly Val Ile Val
 915 920 925
 Gly Asn Asn Pro Asp Leu Lys Leu Asn Asn Asp Lys Thr Ile Thr Leu
 930 935 940
 His Met Gly Lys Ala His Lys Asn Gln Leu Tyr Arg Ala Leu Val Leu
 945 950 955 960
 Ser Asn Asp Ser Gly Ile Asp Val Tyr Asp Ser Asp Asp Lys Ala Pro
 965 970 975
 Thr Leu Arg Thr Asn Asp Asn Gly Asp Leu Ile Phe His Lys Thr Asn
 980 985 990

Thr Phe Val Lys Gln Asp Gly Thr Ile Ile Asn Tyr Glu Met Lys Gly
 995 1000 1005
 Ser Leu Asn Ala Leu Ile Ser Gly Tyr Leu Gly Val Trp Val Pro Val
 1010 1015 1020
 Gly Ala Ser Asp Ser Gln Asp Ala Arg Thr Val Ala Thr Glu Ser Ser
 025 1030 1035 1040
 Ser Ser Asn Asp Gly Ser Val Phe His Ser Asn Ala Ala Leu Asp Ser
 1045 1050 1055
 Asn Val Ile Tyr Glu Gly Phe Ser Asn Phe Gln Ala Met Pro Thr Ser
 1060 1065 1070
 Pro Glu Gln Ser Thr Asn Val Val Ile Ala Thr Lys Ala Asn Leu Phe
 1075 1080 1085
 Lys Glu Leu Gly Ile Thr Ser Phe Glu Leu Ala Pro Gln Tyr Arg Ser
 1090 1095 1100
 Ser Gly Asp Thr Asn Tyr Gly Gly Met Ser Phe Leu Asp Ser Phe Leu
 105 1110 1115 1120
 Asn Asn Gly Tyr Ala Phe Thr Asp Arg Tyr Asp Leu Gly Phe Asn Lys
 1125 1130 1135
 Ala Asp Gly Asn Pro Asn Pro Thr Lys Tyr Gly Thr Asp Gln Asp Leu
 1140 1145 1150
 Arg Asn Ala Ile Glu Ala Leu His Lys Asn Gly Met Gln Ala Ile Ala
 1155 1160 1165
 Asp Trp Val Pro Asp Gln Ile Tyr Ala Leu Pro Gly Lys Glu Val Val
 1170 1175 1180
 Thr Ala Thr Arg Val Asp Glu Arg Gly Asn Gln Leu Lys Asp Thr Asp
 185 1190 1195 1200
 Phe Val Asn Leu Leu Tyr Val Ala Asn Thr Lys Ser Ser Gly Val Asp
 1205 1210 1215
 Tyr Gln Ala Lys Tyr Gly Gly Glu Phe Leu Asp Lys Leu Arg Glu Glu
 1220 1225 1230
 Tyr Pro Ser Leu Phe Lys Gln Asn Gln Val Ser Thr Gly Gln Pro Ile
 1235 1240 1245
 Asp Ala Ser Thr Lys Ile Lys Gln Trp Ser Ala Lys Tyr Met Asn Gly
 1250 1255 1260
 Thr Asn Ile Leu His Arg Gly Ala Tyr Tyr Val Leu Lys Asp Trp Ala
 265 1270 1275 1280
 Thr Asn Gln Tyr Phe Asn Ile Ala Lys Thr Asn Glu Val Phe Leu Pro
 1285 1290 1295
 Leu Gln Leu Gln Asn Lys Asp Ala Gln Thr Gly Phe Ile Ser Asp Ala
 1300 1305 1310

Ser Gly Val Lys Tyr Tyr Ser Ile Ser Gly Tyr Gln Ala Lys Asp Thr
 1315 1320 1325
 Phe Ile Glu Asp Gly Asn Gly Asn Trp Tyr Tyr Phe Asp Lys Asp Gly
 1330 1335 1340
 Tyr Met Val Arg Ser Gln Gln Gly Glu Asn Pro Ile Arg Thr Val Glu
 345 1350 1355 1360
 Thr Ser Val Asn Thr Arg Asn Gly Asn Tyr Tyr Phe Met Pro Asn Gly
 1365 1370 1375
 Val Glu Leu Arg Lys Gly Phe Gly Thr Asp Asn Ser Gly Asn Val Tyr
 1380 1385 1390
 Tyr Phe Asp Asp Gln Gly Lys Met Val Arg Asp Lys Tyr Ile Asn Asp
 1395 1400 1405
 Asp Ala Asn Asn Phe Tyr His Leu Asn Val Asp Gly Thr Met Ser Arg
 1410 1415 1420
 Gly Leu Phe Lys Phe Asp Ser Asp Thr Leu Gln Tyr Phe Ala Ser Asn
 425 1430 1435 1440
 Gly Val Gln Ile Lys Asp Ser Tyr Ala Lys Asp Ser Lys Gly Asn Lys
 1445 1450 1455
 Tyr Tyr Phe Asp Ser Ala Thr Gly Asn Asn Asp Thr Gly Lys Ala Gln
 1460 1465 1470
 Thr Trp Asp Gly Asn Gly Tyr Tyr Ile Thr Ile Asp Ser Asp Ala Asn
 1475 1480 1485
 Asn Thr Ile Gly Val Asn Thr Asp Tyr Thr Ala Tyr Ile Thr Ser Ser
 1490 1495 1500
 Leu Arg Glu Asp Gly Leu Phe Ala Asn Ala Pro Tyr Gly Val Val Thr
 505 1510 1515 1520
 Lys Asp Gln Asn Gly Asn Asp Leu Lys Trp Gln Tyr Ile Asn His Thr
 1525 1530 1535
 Lys Gln Tyr Glu Gly Gln Gln Val Gln Val Thr Arg Gln Tyr Thr Asp
 1540 1545 1550
 Ser Lys Gly Val Ser Trp Asn Leu Ile Thr Phe Ala Gly Gly Asp Leu
 1555 1560 1565
 Gln Gly Gln Arg Leu Trp Val Asp Ser Arg Ala Leu Thr Met Thr Pro
 1570 1575 1580
 Phe Lys Thr Met Asn Gln Ile Ser Phe Ile Ser Tyr Ala Asn Arg Asn
 585 1590 1595 1600
 Asp Gly Leu Phe Leu Asn Ala Pro Tyr Gln Val Lys Gly Tyr Gln Leu
 1605 1610 1615
 Ala Gly Met Ser Asn Gln Tyr Lys Gly Gln Gln Val Thr Ile Ala Gly
 1620 1625 1630

Val Ala Asn Val Ser Gly Lys Asp Trp Ser Leu Ile Ser Phe Asn Gly
 1635 1640 1645
 Thr Gln Tyr Trp Ile Asp Ser Gln Ala Leu Asn Thr Asn Phe Thr His
 1650 1655 1660
 Asp Met Asn Gln Lys Val Phe Val Asn Thr Thr Ser Asn Leu Asp Gly
 665 1670 1675 1680
 Leu Phe Leu Asn Ala Pro Tyr Arg Gln Pro Gly Tyr Lys Leu Ala Gly
 1685 1690 1695
 Leu Ala Lys Asn Tyr Asn Asn Gln Thr Val Thr Val Ser Gln Gln Tyr
 1700 1705 1710
 Phe Asp Asp Gln Gly Thr Val Trp Ser Gln Val Val Leu Gly Gly Gln
 1715 1720 1725
 Thr Val Trp Val Asp Asn His Ala Leu Ala Gln Met Gln Val Ser Asp
 1730 1735 1740
 Thr Asp Gln Gln Leu Tyr Val Asn Ser Asn Gly Arg Asn Asp Gly Leu
 745 1750 1755 1760
 Phe Leu Asn Ala Pro Tyr Arg Gly Gln Gly Ser Gln Leu Ile Gly Met
 1765 1770 1775
 Thr Ala Asp Tyr Asn Gly Gln His Val Gln Val Thr Lys Gln Gly Gln
 1780 1785 1790
 Asp Ala Tyr Gly Ala Gln Trp Arg Leu Ile Thr Leu Asn Asn Gln Gln
 1795 1800 1805
 Val Trp Val Asp Ser Arg Ala Leu Ser Thr Thr Ile Met Gln Ala Met
 1810 1815 1820
 Asn Asp Asn Met Tyr Val Asn Ser Ser Gln Arg Thr Asp Gly Leu Trp
 825 1830 1835 1840
 Leu Asn Ala Pro Tyr Thr Met Ser Gly Ala Lys Trp Ala Gly Asp Thr
 1845 1850 1855
 Arg Ser Ala Asn Gly Arg Tyr Val His Ile Ser Lys Ala Tyr Ser Asn
 1860 1865 1870
 Glu Val Gly Asn Thr Tyr Tyr Leu Thr Asn Leu Asn Gly Gln Ser Thr
 1875 1880 1885
 Trp Ile Asp Lys Arg Ala Phe Thr Val Thr Phe Asp Gln Val Val Ala
 1890 1895 1900
 Leu Asn Ala Thr Ile Val Ala Arg Gln Arg Pro Asp Gly Met Phe Lys
 905 1910 1915 1920
 Thr Ala Pro Tyr Gly Glu Ala Gly Ala Gln Phe Val Asp Tyr Val Thr
 1925 1930 1935
 Asn Tyr Asn Gln Gln Thr Val Pro Val Thr Lys Gln His Ser Asp Ala
 1940 1945 1950

18/36

Gln Gly Asn Gln Trp Tyr Leu Ala Thr Val Asn Gly Thr Gln Tyr Trp
 1955 1960 1965

Ile Asp Gln Arg Ser Phe Ser Pro Val Val Thr Lys Val Val Asp Tyr
 1970 1975 1980

Gln Ala Lys Ile Val Pro Arg Thr Thr Arg Asp Gly Val Phe Ser Gly
 985 1990 1995 2000

Ala Pro Tyr Gly Glu Val Asn Ala Lys Leu Val Asn Met Ala Thr Ala
 2005 2010 2015

Tyr Gln Asn Gln Val Val His Ala Thr Gly Glu Tyr Thr Asn Ala Ser
 2020 2025 2030

Gly Ile Thr Trp Ser Gln Phe Ala Leu Ser Gly Gln Glu Asp Lys Leu
 2035 2040 2045

Trp Ile Asp Lys Arg Ala Leu Gln Ala
 2050 2055

<210> 3
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: artificial
 sequence

<400> 3
 actgcggccg catgccattt acagaaaaag 30

<210> 4
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: artificial
 sequence

<400> 4
 actgctcgag ttatgctgac acagcatttc 30

<210> 5
 <211> 7
 <212> PRT
 <213> Leuconostoc mesenteroides

<400> 5
 Lys Thr Asn Glu Val Phe Leu
 1 5

<210> 6

<211> 7
<212> PRT
<213> Leuconostoc mesenteroides

<400> 6
Lys Phe Asp Ser Asp Arg Leu
1 5

<210> 7
<211> 19
<212> PRT
<213> Leuconostoc mesenteroides

<400> 7
Lys Asn Met Gln Asp Thr Phe Thr Phe Asp Gln Leu Ala Gln Gly Met
1 5 10 15

Glu Phe Tyr

<210> 8
<211> 15
<212> PRT
<213> Leuconostoc mesenteroides

<400> 8
Lys Ser Thr Glu Trp Leu Arg Asp Ala Ile Asp Leu Phe Val Lys
1 5 10 15

<210> 9
<211> 22
<212> PRT
<213> Leuconostoc mesenteroides

<400> 9
Lys Gly Ser Glu Phe Leu Leu Ala Asn Asp Ile Asp Asn Ser Asn Pro
1 5 10 15

Ile Val Gln Ala Glu Gln
20

<210> 10
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 10
ytgrtcraan gtraangtrt cytgcatrtt ytt

<210> 11
<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 11

gaygcnathg ayytnttygt naar

24

<210> 12

<211> 837

<212> DNA

<213> *Leuconostoc mesenteroides*

<400> 12

gatgcaattg	atagttttgt	gaagacgcaa	gctaattgga	ataagcaaac	agaagatgaa	60
gctttcga	gtttgcagtg	gcttcaagg	ggattcctag	cttatcaaga	tgattcacat	120
cggacgccga	atactgattc	aggaaataac	agaaaactag	gacgtcaacc	aattaatatc	180
gatggttcga	aagatacaac	tgatggtaaa	ggctctgaat	tcttattagc	taacgatatt	240
gacaaactcaa	atccgattgt	tcaagctgag	caattaaact	ggctacacta	tttaatgaat	300
tttggtagta	ttacaggtaa	taatgacaat	gcgaattttg	atggcattcg	tgtagatgct	360
gttgataatg	ttgatgctga	tttactaaaa	atagctggcg	attattttaa	agctctatat	420
ggtacagata	aaagcgacgc	caatgccaat	aagcattttg	ctatttttaga	agactggaac	480
ggtaaagatc	ctcagtatgt	taatcaacag	ggcaatgcgc	aattaacaat	ggattacaca	540
gttacttcac	agtttgga	ttctctaaca	catggcgcca	acaacaggag	taacatgtgg	600
tatttcttag	atactggcta	ttatcttaat	ggagatctta	ataagaagat	agtagataag	660
aaccgtccaa	attctggcac	tttggttaac	agaattgcta	attcagggtga	tacaaaagtt	720
attccaaatt	atagttttgt	tagagcacat	gattacgatg	ctcaagatcc	aattagaaaa	780
gccatgattg	atcatggtat	tattaaaaac	atgcaggata	ctttcacttt	tgaccaa	837

<210> 13

<211> 2862

<212> DNA

<213> *Leuconostoc mesenteroides*

<400> 13

gatcaaatag	atgctaccgt	tatacagagac	gttaaggatt	atcccaatct	agtcgttttg	60
cgtaccttgt	cgaaagctta	tggctcggca	aacctgagaa	tcggctatgg	cgctcatgcag	120
gaaccacttt	atcaggttat	gcaggccgta	cgtttaccat	ataattttaa	tacctatcaa	180
atcacagggtg	cagtagctgc	ccttagtgat	caactttatc	tgcaatcagt	tggtgctaag	240
gtgaagtctg	aacgtgaaaa	atttgaacaa	tttttgacga	aacaccagtt	taagtattat	300
caatcacaaa	ccaactttct	ctggattaaa	gttgggtgatg	cgaaacgtgt	tggtgaggct	360
cttctgtcag	aagggatatca	aattaatgac	cgctaaatg	ccgaatggat	tcgcattgca	420
ttaggaactg	tgtctgataa	tgaggggatg	cagcgcattt	tattgaattg	ttaaaaata	480
gctaagagag	tatgttcttc	tcttacctat	ttttatttgt	aattcctatt	atttaatttt	540
gcatgacaat	attaatagcg	tgttacgatt	ctactattta	atgttaataa	aattaataaa	600
tatggtatta	tcttatatgg	gtgatagatg	caccaaatac	tgtatcatgt	ctggtcacat	660
gaaagggaga	ataattaatg	aaacaacaag	aaacagttac	ccgtaaaaaa	ctttataaat	720
ccggtaagggt	ttgggttgca	gcagctactg	catttgcggt	attgggggtt	tcaactgtaa	780
caacagtcca	tgcggataca	aattcgaatg	tcgctgttaa	gcaaataaat	aatacaggaa	840
ccaatgattc	tggcgaaaaa	aaggtaccgg	ttccatcaac	taataatgat	agtttgaagc	900
aaggaacaga	tggtttttgg	tatgattcag	acggcaatcg	tgtcgatcag	aagaccaatc	960
agattctgct	tactcgggaa	caacttaaaa	aaaaataacga	aaaaaattta	tcagtaaatc	1020
gtgatgatac	atcaaaaaaa	gatgatgaaa	atatttctaa	gcagaccaa	attgctaatac	1080
aacaaacagt	agatactgct	aaaggcctga	ctaccagtaa	tttatctgat	cccactactg	1140
ggggctacta	tgaaaatcac	aatggctact	ttgtttatat	agatgcttca	ggaaaacaag	1200
taacagggttt	gcaaaatatt	gatggtaatt	tacaatat	tgatgacaat	ggatatcaag	1260

tcaagggatc	cttccgagat	gtcaacggca	agcatatcta	ttttgattca	gtaacaggga	1320
aagctagttc	aaatgttgat	attgtttaacg	gtaaagctca	aggatatgat	gcgcaaggca	1380
accaattaaa	gaaaagtatt	gtcgccgata	gttctgggca	aacttactat	tttgatggta	1440
atggccaacc	gttaatcggc	ttgcaaaca	ttgatgggaa	cctacaatat	tttaaccaac	1500
aaggggttca	aataaagggt	ggtttccaag	atgttaacaa	taaacgtatt	tattttgcac	1560
caaacacagg	taatgccgtt	gccaatactg	aaataattaa	cggtaaatta	caggggcgtg	1620
acgcaaattg	taaccaggta	aagaatgcat	ttagtaaaga	tgttgcagga	aatacatttt	1680
attttgacgc	aaacggtgtg	atgttaacag	ggttgcaaac	tatttcagga	aagacatatt	1740
atcttgatga	acaaggacac	ctgagaaaaa	attacgcggg	aacattcaat	aatcagttta	1800
tgtacttcga	tgctgataca	gggtcgggta	aaacagcgat	tgaatatcaa	tttgatcaag	1860
gattgggtatc	acaaagtaat	gaaaatactc	ctcacaatgc	cgcaaagtct	tatgataaaa	1920
gtagttttga	aaatgttgat	ggttacttaa	cagcagatac	atggtatcgt	ccaaccgata	1980
ttttaaaaaa	tggagatact	tggacggcat	ctaccgaaac	tgatatgcgt	ccgcttttaa	2040
tgacatgggtg	gcctgacaaa	caaacacaag	caaattactt	gaattttatg	tctagtaaag	2100
gacttggtat	aacgaccact	tatacagcag	ctacgtcaca	aaaaacacta	aatgacgcag	2160
cctttgttat	tcaaacagca	attgaacaac	aaatatcttt	gaaaaaaagt	actgagtggg	2220
tacgtgatgc	aattgatagt	tttgtgaaga	cgcaagctaa	ttggaataag	caaacagaag	2280
atgaagcttt	cgatggtttg	cagtggcttc	aagggggatt	cctagcttat	caagatgatt	2340
cacatcggac	gccgaatact	gattcaggaa	ataacagaaa	actaggacgt	caaccaatta	2400
atatcgatgg	ttcgaaaagt	acaactgatg	gtaaaggctc	tgaattctta	ttagctaacg	2460
atattgacaa	ctcaaatccg	attgttcaag	ctgagcaatt	aaactggcta	cactatttaa	2520
tgaattttgg	tagtattaca	ggtaataatg	acaatgcgaa	ttttgatggc	attcgtgtag	2580
atgctgttga	taatgttgat	gctgatttac	taaaaatagc	tggcgattat	tttaaagctc	2640
tatatggtac	agataaaaagc	gacgccaatg	ccaataagca	tttgtctatt	ttagaagact	2700
ggaacggtaa	agatcctcag	tatgttaatc	aacagggcaa	tgcgcaatta	acaatggatt	2760
acacagttac	ttcacagttt	ggcaattctc	taacacatgg	cgccaacaac	aggagtaaca	2820
tgtgggtat	cttagatact	ggctattatc	ttaatggaga	tc		2862

<210> 14

<211> 3970

<212> DNA

<213> *Leuconostoc mesenteroides*

<400> 14

gatcaaggat	tggtatcaca	aagtaatgaa	aatactcctc	acaatgccgc	aaagtcttat	60
gataaaaagta	gttttgaaaa	tggtgatggg	tacttaacag	cagatacatg	gtatcgtcca	120
accgatattt	taaaaaatgg	agatacttgg	acggcatcta	ccgaaactga	tatgcgtccg	180
cttttaaatga	catggtggcc	tgacaaacaa	acacaagcaa	attacttgaa	tttttgctct	240
agtaaaggac	ttggtataac	gaccacttat	acagcagcta	cgtcacaaaa	aacactaaat	300
gacgcagcct	ttgttattca	aacagcaatt	gaacaacaaa	tatctttgaa	aaaaagtact	360
gagtgggttac	tgatgcaat	tgatagtttt	gtgaagacgc	aagctaattg	gaataagcaa	420
acagaagatg	aagctttcga	tggtttgcag	tggtctcaag	ggggattcct	agcttatcaa	480
gatgattcac	atcggacgcc	gaatactgat	tcaggaaata	acagaaaact	aggacgtcaa	540
ccaattaata	tcgatggttc	gaaagataca	actgatggta	aaggctctga	attcttatta	600
gctaaccgata	ttgacaactc	aaatccgatt	gttcaagctg	agcaattaaa	ctggctacac	660
tattttaatga	attttggtag	tattacaggt	aataatgaca	atgcgaattt	tgatggcatt	720
cgtgtagatg	ctggtgataa	tggtgatgct	gatttactaa	aaatagctgg	cgattatttt	780
aaagctctat	atggtacaga	taaaagcgac	gccaatgccca	ataagcattt	gtctatttta	840
gaagactgga	acggtaaaga	tcctcagtat	gttaatcaac	agggcaatgc	gcaattaaca	900
atggattaca	cagttacttc	acagtttggc	aattctctaa	cacatggcgc	caacaacagg	960
agtaacatgt	ggtatttctt	agatactggc	tattatctta	atggagatct	taataagaag	1020
atagtagata	agaaccgtcc	aaattctggc	actttgggta	acagaattgc	taattcagggt	1080
gatacaaaaag	ttattccaaa	ttatagtttt	gtagagcac	atgattacga	tgctcaagat	1140
ccaattagaa	aagccatgat	tgatcatggt	attattaaaa	acatgcagga	tactttcact	1200
tttgaccaac	tggtctcagg	aatggaaattc	tactataaag	atcaagagaa	tccgtctggt	1260
ttcaaaaagt	ataacgatta	taacttacct	agtgcttatg	caatgttggt	gactaataag	1320
gatactgtac	ctcgtgtcta	ttatggagat	atgtacctcg	aaggcgggca	atatatggaa	1380
aaagggacga	tttacaatcc	tgtcatttca	gcgttgctca	aagctagaat	aaaatatggt	1440
tctgggtgggc	aaacaatggc	taccgatagt	tctggaaaag	accttaaaga	tggcgaaact	1500

```

gattttgttaa caagtgttcg attttggtaaa ggaattatga catcagatca aaccacaaca 1560
caagacaata gccaaagatta taaaaaatcaa ggcacgcgtg tcattgttgg taataaccct 1620
gaccttaagt tgaacaatga taagaccatt accttgcata tgggaaaggc gcataagaat 1680
caactttacc gtgccttagt attatcaaat gactcaggaa ttgatgttta tgatagtgat 1740
gataaagcac caactttgag aacaaatgac aacggtgact tgattttcca taagacaaat 1800
acgtttgtga agcaagatgg aactattata aattacgaaa tgaagggatc attaaatgct 1860
ttaatttcag gttatttagg tgtctgggtg ccagttggag ctagtgattc acaagatgct 1920
cgtacagtg gcaactgagtc atcatcaagt aatgatgggt ctgtattcca ttcaaagtct 1980
gcattagatt ctaatgttat atatgaaggc ttttcaaact ttcaagcgat gccgacttct 2040
cctgagcaaa gtacaaatgt tgttattgca acaaaggcta acttatttaa agaattaggt 2100
attactagtt ttgagttagc acctcaatat aggtctagtg gtgacactaa ttacgggtgc 2160
atgtcattct tagattcttt cttaaataat gggtatgcat ttaccgatag atatgattta 2220
ggctttaaca aagcagacgg gaatcctaac ccaacaaagt atggaacaga tcaagattta 2280
cgtaatgcaa tagaggcatt acacaaaaac ggcacgcagg ctatagctga ttgggttct 2340
gaccaaatat atgctttacc aggaaaggaa gttgttaccg ctactagagt agacgaacgg 2400
ggaaatcaac taaaagacac agattttgtc aacttactct atgttgctaa tactaaaagt 2460
agtgggtgtg attatcaggc aaagtatggc ggcgaatttt tagataaatt aagagaagag 2520
tacctacgt tattcaaaca gaaccaagta tcgacaggtc agccaattga tgcttctaca 2580
aaaattaaag aatggtcagc taaatatatg aatgggacca atattttaca tcgaggtgct 2640
tattatgttt tgaaagactg ggctactaac cagtatttta acattgcaaa aacgaatgaa 2700
gtatttttgc cactacagtt gcagaataaa gatgcgcaaa ctggtttcat tagtgatgcc 2760
tcgggtgtaa aatattactc aattagtggg tatcaagcaa aagatacttt tattgaagat 2820
ggtaatggga attggtatta ctttgataaa gatgggtaca tgggtgcgttc gcagcaagga 2880
gaaaatccta taagaacagt cgaaactagt gtcaacacac gaaacggtaa ttattacttt 2940
atgccaatg gtgtcgagtt gcgcaaaggc tttggaacgg ataatagtgg taatgtctat 3000
tattttgatg atcaaggtaa gatggtgaga gataaataca ttaacgatga tgctaataat 3060
tttttactact taaatgttga tgggactatg tctcgaggac tatttaaatt tgattctgat 3120
actctacagt attttgctag taatgggtgc caaataaaag atagttatgc gaaggatagt 3180
aaaggcaata aatattattt tgactcagct acaggaaata acgatactgg gaaagcccaa 3240
acttgggatg gtaatggcta ctatattact attgattctg atgcgaacaa tacaattggg 3300
gttaacacag actacactgc ctacatcact agctcgctgc gcgaagatgg cttatttget 3360
aacgcacctt acggtgttgt aacaaaagac caaatggta acgatcttaa gtggcagtat 3420
attaaccata cgaaacagta cgaagggcaa caagtgcaag tcacgcgtca atacacagac 3480
agtaaggggag tcagctggaa cttaattacc tttgctgggtg gtgatttaca aggacaaagg 3540
ctttgggtgg atagtcgtgc gttaactatg acaccattta aaacgatgaa ccaaataagc 3600
ttcattagtt atgctaaccg caatgatggg ttgtttttga atgcgccata ccaagtcaag 3660
gggtatcaat tagctgggat gtccaaccaa tacaagggcc aacaagtgc cattgctggg 3720
gtggcgaaag tttctggaaa agactggagt ctgattagtt ttaatgggac acagtactgc 3780
attgtagtgc aggcattgaa taccaatttc acacatgaca tgaaccaaaa ggtctttgtc 3840
aatacaacta gtaatcttga tgggttattc ttaaattgcgc cataccgtca accgggttat 3900
aagttagccg gtttggctaa aaattacaac aaccaaacgg ttactgttag tcaacagtac 3960
tttgatgac 3970

```

<210> 15

<211> 1917

<212> DNA

<213> *Leuconostoc mesenteroides*

<400> 15

```

gatcaaggta agatgggtgag agataaatac attaacgatg atgctaataa tttttatcac 60
ttaaattgtt atgggactat gtctcgagga ctatttaaatt ttgattctga tactctacag 120
tattttgcta gtaatggtgt ccaaataaaa gatagttatg cgaaggatag taaaggcaat 180
aaatattatt ttgactcagc tacaggaaat aacgatactg ggaaagccca aacttgggat 240
ggtaattggct actatattac tattgattct gatgcgaaca atacaattgg ggtaaacaca 300
gactacactg cctacatcac tagctcgctg cgcgaagatg gcttatttgc taacgcacct 360
tacgggtgtt taacaaaaga ccaaattggg aacgatctta agtggcagta tattaacct 420
acgaaacagt acgaagggca acaagtgcaa gtcacgcgtc aatacacaga cagtaaggga 480
gtcagctgga acttaattac ctttgcgtgg ggtgatttac aaggacaaag gctttgggtg 540
gatagtcgtg cgtaactat gacaccattt aaaacgatga accaaataag cttcattagt 600

```

tatgctaacc	gcaatgatgg	gttgtttttg	aatgcgccat	accaagtcaa	ggggtatcaa	660
ttagctggga	tgtccaacca	atacaagggc	caacaagtga	ccattgctgg	ggtggcggaac	720
gtttctggaa	aagactggag	tctgattagt	tttaattggga	cacagtactg	gattgatagt	780
caggcattga	ataccaattt	cacacatgac	atgaaccaaa	aggtctttgt	caatacaact	840
agtaatcttg	atgggttatt	cttaaatgcg	ccataccgtc	aaccgggtta	taagttagcc	900
ggtttggtta	aaaattacaa	caaccaaacg	gttactgtta	gtcaacagta	ctttgatgat	960
caaggcacgg	tctggagtca	ggttgtcctt	gggggtcaga	cggtctgggt	tgataaccat	1020
gcattggcac	agatgcaagt	tagtgataca	gaccaacagc	tctatgtgaa	tagcaatggg	1080
cggaatgatg	ggttattctt	gaatgcgcca	tatcgtgggc	aagggtcaca	actgataggc	1140
atgacggcag	attataatgg	gcaacatgta	caagtgaacca	agcaagggca	agatgcctat	1200
ggtgcacaa	ggcgtcttat	tacgctaaat	aatcaacagg	tctgggttga	tagtcgcgct	1260
ttgagcacaa	caatcatgca	agccatgaat	gataatatgt	atgtaaatag	cagccaacgg	1320
acagatggct	tgtggttaaa	cgcaccttat	acgatgagtg	gggctaaatg	ggctgggtgat	1380
acacgttcag	ctaattggcg	ctatgtccat	atttcaaaaag	cttattcaaa	cgaagtcggc	1440
aatacatatt	acttgacgaa	tttgaatggg	caaagcacat	ggattgacaa	gcgggcggtt	1500
actgtgacct	tcgatcaggt	ggtggcatta	aatgcaacga	ttgtggcacg	ccaacgacca	1560
gatgggatgt	ttaagacagc	accatatggg	gaagcggggg	cgcagtttgt	cgattatgtg	1620
acaaactata	accagcaaac	cgtgccagta	acaaagcaac	attcagatgc	tcaggggaat	1680
caatggtact	tagcgacagt	gaatgggaca	caatactgga	ttgatcaacg	gtcattttca	1740
ccagtagtaa	cgaaggtggg	tgattatcaa	gctaagattg	tgccacggac	aacacgtgat	1800
ggtgtgttta	gtggcgccac	ctatggggaa	gtgaatgcta	agctagttaa	catggcaact	1860
gcgtatcaaa	atcaagttgt	ccatgcgaca	ggggaatata	cgaatgcttc	agggatc	1917

<210> 16

<211> 4066

<212> DNA

<213> *Leuconostoc mesenteroides*

<400> 16

gatcttaagt	ggcagtatat	taaccatacg	aaacagtacg	aaggggcaaca	agtgcaagtc	60
acgcgtcaat	acacagacag	taagggagtc	agctggaact	taattacctt	tgctgggtgg	120
gatttacaag	gacaaaggct	ttgggtggat	agtcgtgcgt	taactatgac	accattttaa	180
acgatgaacc	aaataagctt	cattagttat	gctaaccgca	atgatgggtt	gtttttgaat	240
gcgccatacc	aagtcaaggg	gtatcaatta	gctgggatgt	ccaaccaata	caagggccaa	300
caagtgaacca	ttgctggggg	ggcgaacggt	tctggaaaag	actggagtc	gatttagttt	360
aatgggacac	agtactggat	tgatagtcag	gcattgaata	ccaatttcac	acatgacatg	420
aaccaaaggg	tctttgtcaa	tacaactagt	aatcttgatg	ggttattctt	aaatgcgcca	480
taccgtcaac	cgggttataa	gtagccgggt	ttggctaaaa	attacaacaa	ccaaacgggt	540
actgttagtc	aacgatctt	tgatgatcaa	ggcacggtct	ggagtcaggt	tgctcttggt	600
ggtcagacgg	tctgggttga	taaccatgca	ttggcacaga	tgcaagttag	tgatacagac	660
caacagctct	atgtgaatag	caatggctcg	aatgatgggt	tattcttgaa	tgccgccatat	720
cgtggtcaag	ggtcacaact	gataggcatg	acggcagatt	ataatgggca	acatgtacaa	780
gtgaccaagc	aaggggcaaga	tgccataggt	gcacaatggc	gtcttattac	gctaaataat	840
caacaggtct	gggttgatag	tcgcgccttg	agcacaacaa	tcagtgaagc	catgaatgat	900
aatatgtatg	taaatagcag	ccaacggaca	gatggcttgt	ggttaaagcg	accttatacg	960
atgagtgggg	ctaaatgggc	tggtgataca	cgttcagcta	atgggcgcta	tgcccatatt	1020
tcaaaagctt	attcaaacga	agtcggcaat	acatattact	tgacgaattt	gaatgggtcaa	1080
agcacatgga	ttgacaagcg	ggcgtttact	gtgaccttcg	atcaggtggg	ggcattaaat	1140
gcaacgattg	tggcacgcca	acgaccagat	gggatgttta	agacagcacc	atatggtgaa	1200
gcggggggcg	agtttgtcga	ttatgtgaca	aactataacc	agcaaaccgt	gccagtaaca	1260
aagcaacatt	cagatgctca	ggggaatcaa	tggtacttag	cgacagtga	tgggacacaa	1320
tactggattg	atcaacggtc	attttcacca	gtagtaacga	aggtgggtga	ttatcaagct	1380
aagattgtgc	cacggacaac	acgtgatggg	gtgttttagt	gcgcacccta	tggggaagtg	1440
aatgctaagc	tagttaacat	ggcaactgcg	tatcaaaatc	aagttgtcca	tgcgacaggg	1500
gaatatacga	atgcttcagg	gatcacatgt	agtcagttcg	cgttaagcgg	gcaagaagac	1560
aagctatgga	ttgataagcg	tgctttgcaa	gcttaaggga	aggattcgac	aaaggagggt	1620
aacattatca	gcggatgggt	ttatctctct	ttcctgtact	cagtatttcc	caaataattg	1680
agacagtttc	atgacaaaac	aacaaaacta	gtgtcaatgc	ctcgggttat	gggtaaaacta	1740
ctattagtta	aagggttggg	gcataataat	atcacattca	atatattatg	tattttttat	1800

tgattatgtg	atTTTTtGag	atttggagcg	aaaatgaata	aagtaataat	taataaaaagt	1860
ttctgtgttt	tggtaatgag	tttgttgtcg	atatttctat	tctcattaag	agtagacgct	1920
cgttccaata	gtggctacaa	tcaaatatct	tttaatcaac	atgaaattgc	ttatgcgct	1980
tttagtcaag	taccttggtg	tgcaactgtg	aatgtcggaa	tgtctgcaga	caaaaataat	2040
atctatacag	ctattgatat	ggggccatag	tatcaaggaa	cttctttttt	tccgtggtgg	2100
tatcaacatg	ataattatga	ttatcatgat	ggttatcaac	ttgcgattga	caatcaatcg	2160
cttttcatag	aaatggttga	tacaggaagg	tcgtcgtttg	atgttattga	gttgataaca	2220
ggtaaaaaag	ttggttatgg	aaaattcaca	gcctataact	ataacaagac	ttcatggcag	2280
tttcaaatat	ctttaaaggc	gataaagaaa	attttaaata	ataacatatc	agaaaatgcy	2340
acaattagtt	tgtacaatgg	ttatctattc	aaacaagccg	aaagtatcac	gtatgctggt	2400
tcaccaacag	ggccatgggt	acttgccggt	attggtttgg	tttttgctgg	tgggggctat	2460
tacctgcaaa	atcagcgccg	aaaaacgatt	acggcatcat	caacaaacga	ggtaacccat	2520
gcttaataat	gctgttatta	ttgggtctaat	tattatttgg	gtctatgtat	tatctgttct	2580
gaagcgtgct	aaaacggatg	cgttttattt	tttggctggc	agcgcaggct	tgttcgtcat	2640
tcttctttta	tggagtaaac	cttatgggtg	ttggttgttt	tcgaccattc	tcacttggtc	2700
ggccgggggt	gttgggtcatc	tcactggctt	at ttgatact	ttctatgcgt	cacatgttat	2760
tcaggttgtg	gcgaatcatc	ataccagcat	tttattgggt	gattatgaat	gttcaggcat	2820
tattgaaaca	acggcatttt	ggggattaat	tgccttttat	cctgtgtatg	atgcacaaaa	2880
gcggttaatg	ttggcgctta	ttggggcatt	atggtttttc	cttgcaaata	cgttacgact	2940
atcttttggg	gcagttgtaa	tttattattt	tggcgatggc	gctttttatg	ttgctcattc	3000
aattatcgga	cgccgtttgt	tctattcaat	tgtaatcttg	ctttattatc	tagtcttcat	3060
aaaggggcaa	cttgttaatc	aaatgctcaa	aaaaacagga	taaactatgt	cattttattt	3120
acatcaattt	acttatcagt	taggtttttg	gttcacttgg	ttgctcattc	cgtttgttgt	3180
ggaaattttc	ccagctatta	tttcaatcta	gttttgataa	aaacatcaaa	aaagcatcaa	3240
gtgatgcaag	aaccattgaa	attaccaatg	gtatctattg	tattacctat	atataattca	3300
ggacaaacat	tataccagtg	tattcagtcg	attagtcaat	caacctatcc	aaaacaatta	3360
atccaaatta	ttgccgtgaa	taatcaaagc	actgataaca	gttttacagt	gtttaatcag	3420
gctcaagctg	at ttcccaat	acttagaatg	cagtggatga	acacagatca	aggtaaagcg	3480
cgtgcactaa	atgctgctat	ttataatagt	atggggcaat	atattattaa	tttgatact	3540
gatggttggg	tagaacccaa	tgccttaaaa	cggtttgtgc	tttattttga	aaatcactca	3600
gaaattgatg	tagcaactgg	taogattctg	acacaaaaaa	aaatgattca	aaaaacgcaa	3660
agtaaatggc	ttaaattgct	acagttaaac	gaatattttg	agtatgcaca	atcattctta	3720
tcaggccgta	gtattgaaaa	cogtggtaat	cgtttgttca	caatgtcagg	tgcattttca	3780
gcgttttagac	gtgatgtatt	agttcagaca	tttatgtata	atgttgacac	ggttggtgaa	3840
gatactgata	tgacatttca	actgcgattc	cgcctaggca	agcgtattgg	tttctgtgat	3900
gatgccatgt	tttatgttga	accaatatca	ggttatagtg	aactttattt	acaacggcag	3960
cgttggcagc	gtgggcaaat	tgaagtcgca	caaaatttca	tgcagaataa	gttgagtgtc	4020
cgtcagattt	ttactaactt	tatgattagt	cgattaatga	ttgatc		4066

<210> 17

<211> 7387

<212> DNA

<213> *Leuconostoc mesenteroides*

<400> 17

gatcaaatag	atgctaccgt	tatacgagac	gttaaggatt	atcccaatct	agtcgtttttg	60
cgtaccttgt	cgaaagctta	tggctcggca	aacctgagaa	tcggctatgg	cgtcatgcag	120
gaaccacttt	atcaggttat	gcaggccgta	cgtttaccat	ataattttaa	tacctatcaa	180
atcacagggtg	cagtagctgc	ccttagtgat	caactttatc	tgcaatcagt	tgttgctaag	240
gtgaagtctg	aacgtgaaaa	at ttgaacaa	tttttgacga	aacaccagtt	taagtattat	300
caatcacaaa	ccaactttct	ctggattaaa	gttgggtgatg	cgaaacgtgt	tggtgaggct	360
cttctgtcag	aagggtatca	aattaatgac	cgcctaaatg	ccgaatggat	tcgcattgca	420
ttaggaactg	tgtctgataa	tgaggggatg	cagcgcattt	tattgaattg	ttaaaaata	480
gctaagagag	tatgttcttc	tcttacctat	ttttatttgt	aattcctatt	at ttaatttt	540
gcatgacaat	attaatagcg	tgttacgatt	ctactattta	atgttaataa	aattaataaa	600
tatgggtatta	tcttatatgg	gtgatagatg	caccaaatatc	tgtatcatgt	ctggtcacat	660
gaaagggaga	ataattaatg	aaacaacaag	aaacagttac	ccgtaaaaaa	ctttataaat	720
ccggttaagg	ttgggttgca	gcagctactg	catttgcggt	attgggggtt	tcaactgtaa	780
caacagtcca	tgcggataca	aattcgaatg	tcgctgttaa	gcaaataaat	aatacaggaa	840

ccaatgattc	tggcgaaaaa	aaggtagcgg	ttccatcaac	taataatgat	agtttgaagc	900
aaggaacaga	tgggttttgg	tatgattcag	acggcaatcg	tgtcgatcag	aagaccaatc	960
agattctgct	tactgcggaa	caacttaaaa	aaaataacga	aaaaaattta	tcagtaatca	1020
gtgatgatac	atcaaaaaaa	gatgatgaaa	atattttctaa	gcagacccaaa	attgctaate	1080
aacaaacagt	agatactgct	aaaggcctga	ctaccagtaa	tttatctgat	cccatcactg	1140
ggggctacta	tgaaaatcac	aatggctact	ttgtttatat	agatgcttca	ggaaaacaag	1200
taacagggtt	gcaaaaatatt	gatggtaatt	tacaatatatt	tgatgacaat	ggatatcaag	1260
tcaagggatc	cttccgagat	gtcaacggca	agcatatcta	ttttgattca	gtaacaggga	1320
aagctagttc	aatggttgat	attgttaacg	gtaaagctca	aggatatgat	gcgcaaggca	1380
accaattaaa	gaaaagttaa	gtcgccgata	gttctgggca	aacttactat	tttgatggta	1440
atggccaacc	gttaatcggc	ttgcaaacaa	ttgatgggaa	cctacaatat	tttaaccaac	1500
aaggggttca	aataaagggt	ggtttccaag	atgttaacaa	taaacgtatt	tattttgcac	1560
caaacacagg	taatgccgtt	gccaatactg	aaataattaa	cggtaaatta	caggggcgtg	1620
acgcaaattg	taaccaggta	aagaatgcat	ttagtaaaga	tggtgcagga	aatacatttt	1680
attttgacgc	aaacggtgtg	atgttaacag	ggttgcaaac	tatttcagga	aagacataat	1740
atcttgatga	acaaggacac	ctgagaaaaa	attacgcggg	aacattcaat	aatcagttta	1800
tgtacttcga	tgctgatata	ggtgcgggta	aaacagcgat	tgaatatcaa	tttgatcaag	1860
gatttggtatc	acaaagtaat	gaaaatactc	ctcacaatgc	cgcaaagtct	tatgataaaa	1920
gttagtttga	aatggttgat	ggttacttaa	cagcagatac	atggtatcgt	ccaaccgata	1980
ttttaaaaaa	tgagataact	tgacgcggat	ctaccgaaac	tgatatgcgt	ccgcttttaa	2040
tgacatgggtg	gcctgacaaa	caaacacaag	caaattactt	gaattttatg	tctagtaaag	2100
gacttggtat	aacgaccact	tatacagcag	ctacgtcaca	aaaaacacta	aatgacgcag	2160
cctttgttat	tcaaacagca	attgaacaac	aaatatcttt	gaaaaaaagt	actgagtggg	2220
tacgtgatgc	aattgatagt	tttgtgaaga	cgcaagctaa	ttggaataag	caaacagaag	2280
atgaagcttt	cgatggtttg	cagtggcttc	aagggggatt	cctagcttat	caagatgatt	2340
cacatcggac	gccgaataact	gattcaggaa	ataacagaaa	actaggacgt	caaccaatta	2400
atatcgatgg	ttcgaaagat	acaactgatg	gtaaaggctc	tgaattctta	ttagctaacg	2460
atatggacaa	ctcaaatccg	attgttcaag	ctgagcaatt	aaactggcta	cactatttaa	2520
tgaattttgg	tagtattaca	ggtaataatg	acaatgcgaa	ttttgatggc	attcgtgtag	2580
atgctgttga	taatgttgat	gctgatttac	taaaaatagc	tggcgattat	tttaaagctc	2640
tatatggtac	agataaaaagc	gacgccaatg	ccaataagca	tttgtctatt	ttagaagact	2700
ggaacggtaa	agatcctcag	tatgttaatc	aacaggggcaa	tgcgcaatta	acaatggatt	2760
acacagttac	ttcacagttt	ggcaattctc	taacacatgg	cgccaacaac	aggagtaaca	2820
tgtggtatatt	cttagataact	ggctattatc	ttaatggaga	tcttaataag	aagatagtag	2880
ataagaaccg	tccaaattct	ggcacttttg	ttaacagaat	tgctaattca	ggtgatacaa	2940
aagttattcc	aaattatagt	tttgtttagag	cacatgatta	cgatgctcaa	gatccaatta	3000
gaaaagccat	gattgatcat	ggtattatta	aaaacatgca	ggatactttc	acttttgacc	3060
aactggctca	gggaatggaa	ttctactata	aagatcaaga	gaatccgtct	ggtttcaaaa	3120
agtataacga	ttataactta	cctagtgcct	atgcaatggt	gttgactaat	aaggatactg	3180
tacctcgtgt	ctattatgga	gatatgtacc	tcgaaggcgg	gcaatatatg	gaaaaaggga	3240
cgattttacaa	tcctgtcatt	tcagcgttgc	tcaaagctag	aataaaaatat	gtttctgggtg	3300
ggcaaacaat	ggctaccgat	agttctggaa	aagaccttaa	agatggcgaa	actgatttgt	3360
taacaagtgt	tcgattttgt	aaaggaatta	tgacatcaga	tcaaaccaca	acacaagaca	3420
atagccaaga	ttataaaaaat	caaggcatcg	gtgtcattgt	tggtataaac	cctgacctta	3480
agttgaacaa	tgataagacc	attaccttgc	atatgggaaa	ggcgcataag	aatcaacttt	3540
accgtgcctt	agtattatca	aatgactcag	gaattgatgt	ttatgatagt	gatgataaag	3600
caccaacttt	gagaacaaat	gacaacgggt	acttgatttt	ccataagaca	aatacgtttg	3660
tgaagcaaga	tggaaactatt	ataaattacg	aaatgaaggg	atcattaaat	gctttaattt	3720
caggttatatt	aggtgtctgg	gtgccagttg	gagctagtga	ttcacaagat	gctcgtacag	3780
tggcaactga	gtcatcatca	agtaatgatg	gttctgtatt	ccattcaaatt	gctgcattag	3840
attctaattgt	tatatatgaa	ggcttttcaa	actttcaagc	gatgccgact	tctcctgagc	3900
aaagtacaaa	tggtgttatt	gcaacaaagg	ctaacttatt	taaagaatta	ggtattacta	3960
gttttgagtt	agcacctcaa	tataggtcta	gtgggtgacac	taattacggt	ggcatgtcat	4020
tcttagattc	tttcttaaatt	aatggttatg	cattttaccga	tagatatgat	ttaggcttta	4080
acaaagcaga	cgggaatcct	aacccaacaa	agtgatggaac	agatcaagat	ttacgtaatg	4140
caatagaggc	attacacaaa	aacggcatgc	aggctatagc	tgattgggtt	cctgacccaa	4200
tatatgcttt	accagaaaag	gaagttgtta	ccgctactag	agtagacgaa	cggggaaatc	4260
aactaaaaga	cacagatttt	gtcaacttac	tctatgttgc	taataactaaa	agtagtgggtg	4320
tggattatca	ggcaaagtat	ggcggcgaat	ttttagataa	attaagagaa	gagtacccat	4380
cgttatttcaa	acagaaccaa	gtatcgacag	gtcagccaat	tgatgcttct	acaaaaatta	4440

```

agcaatgggtc agctaaatat atgaatggga ccaatatattt acatcgaggt gcttattatg 4500
ttttgaaaga ctgggctact aaccagtatt ttaacattgc aaaaacgaat gaagtatttt 4560
tgccactaca gtgcagaat aaagatgcgc aaactggttt cattagtgat gcctccggtg 4620
taaaatatata ctcaattagt gggtatcaag caaaagatac ttttattgaa gatggtaatg 4680
ggaattggta ttactttgat aaagatgggt acatgggtgcg ttgcgagcaa ggagaaaatc 4740
ctataagaac agtcgaaact agtgtaaca caccgaaacgg taattattac tttatgccaa 4800
atgggtgtcga gttgcgcaaa ggctttggaa cggataatag tggtaatgtc tattattttg 4860
atgatcaagg taagatgggt agagataaat acattaacga tgatgctaata aattttttatc 4920
acttaaatgt tgatgggact atgtctcgag gactatttaa atttgattct gatactctac 4980
agtattttgc tagtaatggg gtccaaataa aagatagtta tgcgaaggat agtaaaggca 5040
ataaatatta ttttgactca gctacaggaa ataacgatac tgggaaagcc caaacttggg 5100
atggtaatgg ctactatatt actattgatt ctgatgcgaa caatacaatt ggggttaaca 5160
cagactacac tgctacatc actagctcgc tgcgcgaaga tggcttattt gctaacgcac 5220
cttacggtgt tgtaacaaaa gaccaaattg gtaacgatct taagtggcag tatattaacc 5280
atacgaaaca gtacgaaggg caacaagtgc aagtcacgcg tcaatacaca gacagtaagg 5340
gagtcagctg gaacttaatt acctttgctg gtgggtgattt acaaggacaa aggctttggg 5400
tggatagtcg tgcgttaact atgacaccat ttaaaacgat gaaccaataa agcttcatta 5460
gttatgctaa ccgcaatgat ggggtgtttt tgaatgcgcc ataccaagtc aaggggtatc 5520
aattagctgg gatgtccaac caatacaagg gccacaagtg gaccattgct ggggtggcga 5580
acgtttctgg aaaagactgg agtctgatta gttttaatgg gacacagtac tggattgata 5640
gtcaggcatt gaataccaat ttcacacatg acatgaacca aaaggtcttt gtcaatacaa 5700
ctagtaatat tgatgggtta ttcttaaatg cgccataccg tcaaccgggt tataagttag 5760
ccggtttggc taaaaattac aacaaccaa cggttactgt tagtcaacag tactttgatg 5820
atcaaggcac ggtctggagt cagggtgtcc ttgggggtca gacggtctgg gttgataacc 5880
atgcattggc acagatgcaa gttagtata cagaccaaca gctctatgtg aatagcaatg 5940
gtcggaaatga tgggttattc ttgaatgcgc catatcgtgg tcaaggtca caactgatag 6000
gcatgacggc agattataat gggcaacatg tacaagtgc caagcaaggg caagatgcct 6060
atgggtgcaca atggcgctctt attacgctaa ataatcaaca ggtctgggtt gatagtcgcg 6120
ctttgagcac aacaatcatg caagccatga atgataatat gtatgtaaat agcagccaac 6180
ggacagatgg cttgtgggta aacgcacctt atacgatgag tggggctaaa tgggctgggtg 6240
atacacgttc agctaattgg cgctatgtcc atatttcaaa agcttattca aacgaagtgc 6300
gcaatacata ttacttgacg aatttgaatg gtcaaagcac atggattgac aagcgggcgt 6360
ttactgtgac cttcgatcag gtgggtggcat taaatgcaac gattgtggca cgccaacgac 6420
cagatgggat gtttaagaca gcaccatatg gtgaagcggg ggcgcagttt gtogattatg 6480
tgacaaaacta taaccagcaa accgtgccag taacaaagca acattcagat gctcagggga 6540
atcaatggta cttagcgaca gtgaatggga cacaatactg gattgatcaa cggtcatttt 6600
caccagtagt aacgaagggt gttgattatc aagctaagat tgtgccacgg acaacacgtg 6660
atgggtgtgt tagtggcgca ccctatggg aagtgaatgc taagctagtt aacatggcaa 6720
ctgcgtatca aaatcaagtt gtccatgcga cagggaata tacgaatgct tcagggatca 6780
catggagtca gttcgcgtta agcgggcaag aagacaagct atggattgat aagcgtgctt 6840
tgcaagctta agggaaggat tcgacaaagg agggtaacat tatcagcgga tgggtgttatc 6900
ctcctttcct gtactcagta ttcccaaat aattgagaca gtttcatgac aaatcaacaa 6960
aactagtgtc aatgcctcgg ttatggggta aactactatt agttaaaggg ttgttgcata 7020
ataatatcac attcaatata ttatgtattt ttatctgatt atgtgatttt ttgagatttg 7080
gagcgaaaat gaataaagta ataattaata aaagtttctg tgttttggtat agagtttgt 7140
tgtcgatatt tctatttctca ttaagagtag acgctcgttc caatagtggc tacaatcaaa 7200
tatcttttaa tcaacatgaa attgcttatg cgccttttag tcaagtacct tgggatgcaa 7260
ctgtgaatgt cggaatgtct gcagacaaaa ataatatcta tacagctatt gatatggggc 7320
catagtatca aggaacttct ttttttccgt ggtggtatca acatgataat tatgattatc 7380
atgatgg 7387

```

<210> 18

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 18

gggcccgccta gcatgaaaca acaagaaaca gt

32

<210> 19

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 19

cccggggtcg acctttgtcg aatccttccc

30

<210> 20

<211> 6204

<212> DNA

<213> Leuconostoc mesenteroides

<400> 20

```

gctagcatga aacaacaaga aacagttacc cgtaaaaaac tttataaato cggtaaagggtt 60
tggtgtgtag cagctactgc atttgcggta ttggggggtt caactgtaac aacagtccat 120
gcggatacaa attcgaatgt cgctgttaag caaataaata atacaggaac caatgattct 180
ggcgaaaaaa aggtaccggt tccatcaact aataatgata gtttgaagca aggaacagat 240
ggttttttgt atgattcaga cggcaatcgt gtcgatcaga agaccaatca gattctgctt 300
actgcggaac aacttaaaaa aaataacgaa aaaaatttat cagtaatcag tgatgataca 360
tcaaaaaaag atgatgaaaa tattttctaag cagacccaaa ttgctaataca acaaacagta 420
gatactgcta aaggcctgac taccagtaat ttatctgatc ccatcactgg gggtcactat 480
gaaaatcaca atggctactt tgtttatata gatgcttcag gaaaacaagt aacagggtttg 540
caaaatattg atggtaattt acaatatatt gatgacaatg gatatcaagt caagggatcc 600
ttccgagatg tcaacggcaa gcatatctat ttgattcag taacagggaag agctagttca 660
aatgttgata ttgttaacgg taaagctcaa ggatatgatg cgcaaggcaa ccaattaaag 720
aaaagttatg tcgccgatag ttctgggcaa acttactatt ttgatggtaa tggccaaccg 780
ttaatcgggt tgcaacaact tgatgggaac ctacaatatt ttaaccaaca aggggttcaa 840
ataaagggtg gtttccaaga tgtaacaact aaacgtatatt attttgcacc aaacacaggt 900
aatgccgttg ccaatactga aataattaac ggtaaattac aggggcgtga cgcaaatggt 960
aaccaggtaa agaatgcatt tagtaaagat gttgcaggaa atacatttta ttttgacgca 1020
aacggtgtga tgtaaacagg gttgcaaaact atttcaggaa agacatatta tcttgatgaa 1080
caaggacacc tgagaaaaaa ttacgcggga acattcaata atcagtttat gtacttcgat 1140
gctgatacag gtgcgggtaa aacagcgatt gaatatcaat ttgatcaagg attggtatca 1200
caaagtaatg aaaatactcc tcacaatgcc gcaaagtctt atgataaaag tagttttgaa 1260
aatgttgatg gttacttaac agcagatata tggatcgtc caaccgatat tttaaaaaat 1320
ggagatactt ggacggcacc taccgaaact gatatgcgtc cgcttttaac gacatgggtg 1380
cctgacaaac aaacacaagc aaattacttg aattttatgt ctagttaaagg acttgggtata 1440
acgaccactt atacagcagc tacgtcacia aaaacactaa atgacgcagc ctttggttatt 1500
caaacagcaa ttgaacaaca aatatctttg aaaaaaagta ctgagtgggt acgtgatgca 1560
attgatagtt ttgtgaagac gcaagctaatt tgggaataagc aaacagaaga tgaagctttc 1620
gatggtttgc agtggcttca agggggattc cttagcttatc aagatgattc acatcggaag 1680
ccgaatactg attcaggaaa taacagaaaa ctaggacgtc aaccaattaa tatcgatggt 1740
tcgaaagata caactgatgg taaaggctct gaattcttat tagctaacga tattgacaac 1800
tcaaatccga ttgttcaagc tgagcaatta aactggctac actattttaa gaattttggt 1860
agtattacag gtaataatga caatgcgaat tttgatggca ttcgtgtaga tgctgttgat 1920
aatgttgatg ctgatttact aaaaatagct ggcgattatt ttaaagctct atatgggtaca 1980
gataaaaagc acgccaatgc caataagcat ttgtctatatt tagaagactg gaacggtaaa 2040
gatcctcagt atgttaatac acagggcaat gcgcaattaa caatggatta cacagttact 2100
tcacagtttg gcaattctct aacacatggc gccacaaca ggagtaacat gtgggtatttc 2160

```

ttagatactg	gctattatct	taatggagat	cttaataaga	agatagtaga	taagaaccgt	2220
ccaaattctg	gcactttggt	taacagaatt	gctaattcag	gtgatacaaa	agttattcca	2280
aattatagtt	ttgttagagc	acatgattac	gatgctcaag	atccaattag	aaaagccatg	2340
attgatcatg	gtattattaa	aaacatgcag	gatactttca	cttttgacca	actggctcag	2400
ggaatggaat	tctactataa	agatcaagag	aatccgtctg	gtttcaaaaa	gtataacgat	2460
tataacttac	ctagtgttta	tgcaatgttg	ttgactaata	aggatactgt	acctcgtgtc	2520
tattatggag	atatgtacct	cgaaggcggg	caatatatgg	aaaaagggac	gatttacaat	2580
cctgtcattt	cagcgttgct	caaagctaga	ataaaatatg	tttctgggtg	gcaaacaatg	2640
gctaccgata	gttctggaaa	agaccttaaa	gatggcgaaa	ctgattttgt	aacaagtgtt	2700
cgattttgga	aaggaattat	gacatcagat	caaaccacaa	cacaagacaa	tagccaagat	2760
tataaaaaatc	aaggcatcgg	tgtcattggt	ggtaataaacc	ctgaccttaa	gttgaacaa	2820
gataaagacca	ttaccttgca	tatgggaaag	gcgcataaga	atcaacttta	ccgtgcctta	2880
gtattatcaa	atgactcagg	aattgatgtt	tatgatagtg	atgataaagc	accaactttg	2940
agaacaaatg	acaacggtga	cttgattttc	cataagacaa	atacgtttgt	gaagcaagat	3000
ggaactatta	taaattacga	aatgaaggga	tcattaaatg	ctttaatttc	aggttattta	3060
ggtgtctggg	tgccagttgg	agctagtgat	tcacaagatg	ctcgtacagt	ggcaactgag	3120
tcatcatcaa	gtaatgatgg	ttctgtattc	cattcaaatg	ctgcattaga	ttctaagtgt	3180
atatatgaag	gcttttcaaa	ctttcaagcg	atgccgactt	ctcctgagca	aagtacaaat	3240
gttgtttattg	caacaaaggc	taacttattt	aaagaattag	gtattactag	ttttgagtta	3300
gccttcaat	ataggtctag	tggtgacact	aattacgggtg	gcatgtcatt	cttagattct	3360
ttcttaataa	atggttatgc	atttaccgat	agatatgatt	taggctttta	caaagcagac	3420
gggaatccta	acccaacaaa	gtatggaaca	gatcaagatt	tacgtaatgc	aatagaggca	3480
ttacacaaaa	acggcatgca	ggctatagct	gattgggttc	ctgaccaa	atatgcttta	3540
ccaggaaagg	aagttgttac	cgctactaga	gtagacgaac	ggggaaatca	actaaaagac	3600
acagattttg	tcaacttact	ctatgttgct	aatactaaaa	gtagtgggtg	ggattatcag	3660
gcaaaagtatg	gcggcgcaatt	tttagataaa	ttaagagaag	agtaccatc	gttattcaaa	3720
cagaaccaag	tatcgacagg	tcagccaatt	gatgcttcta	caaaaattaa	gcaatgggtca	3780
gctaaatata	tgaatgggac	caatatttta	catcgagggtg	cttattatgt	tttgaaagac	3840
tgggctacta	accagtat	taacattgca	aaaacgaatg	aagtattttt	gccactacag	3900
ttgcagataa	aagatgcgca	aactgggttc	attagtgtg	cctccgggtg	aaaattattac	3960
tcaattagt	gttatcaagc	aaaagatact	tttattgaag	atggtaaatg	gaattgggtat	4020
tactttgata	aagatgggtta	catgggtgct	tcgcagcaag	gagaaaatcc	tataagaaca	4080
gtcgaaacta	gtgtcaacac	acgaaacggg	aattattact	ttatgccaaa	tggtgtcgag	4140
ttgcgcaaag	gctttggaac	ggataatagt	ggtaatgtct	attattttga	tgatcaaggt	4200
aagatgggtga	gagataaata	cattaacgat	gatgctaata	atttttatca	cttaaatgtt	4260
gatgggacta	tgtctcgagg	actattttaa	tttgattctg	atactctaca	gtattttgct	4320
agtaatgggtg	tccaaataaa	agatagttat	gcgaaggata	gtaaaggcaa	taaatattat	4380
tttgactcag	ctacaggaaa	taacgatact	gggaaagccc	aaacttgga	tggtaatggc	4440
tactatat	ctattgattc	tgatgcgaac	aatacaattg	gggttaacac	agactacact	4500
gcctacatca	ctagctcgct	gcgcgaagat	ggcttatttg	ctaacgcacc	ttacgggtgt	4560
gtaacaaaag	accaaagtgg	taacgatctt	aagtggcagt	atattaacca	tacgaaacag	4620
tacgaagggc	aacaagtgca	agtcacgcgt	caatacacag	acagtaaggg	agtcagctgg	4680
aacttaatta	cctttgctgg	tggtgattta	caaggacaaa	ggctttgggt	ggatagtcgt	4740
gcgttaacta	tgacaccatt	taaaacgatg	aaccaataaa	gcttcattag	ttatgctaac	4800
cgcaatgatg	ggttggtttt	gaatgcgcca	taccaagtca	aggggtatca	attagctggg	4860
atgtccaacc	aatacaaggg	ccaacaagtg	accattgctg	gggtggcgaa	cgtttctgga	4920
aaagactgga	gtctgattag	ttttaatggg	acacagtact	ggattgatag	tcaggcattg	4980
aataccaatt	tcacacatga	catgaaccaa	aaggtccttg	tcaatacaac	tagtaattct	5040
gatgggttat	tcttaaatgc	gccataccgt	caaccgggtt	ataagttagc	cggtttggt	5100
aaaaattaca	acaaccaa	ggttactgtt	agtcaacagt	actttgatga	tcaaggcacg	5160
gtctggagtc	aggttgtcct	tgggggtcag	acgggtctgg	ttgataacca	tgcattggca	5220
cagatgcaag	ttagtgtac	agaccaacag	ctctatgtga	atagcaatgg	tcggaatgat	5280
gggttattct	tgaatgcgcc	atctcgtgg	caagggtcac	aactgatagg	catgacggca	5340
gattataatg	ggcaacatgt	acaagtgacc	aagcaagggc	aagatgccta	tggtgcacaa	5400
tggcgtctta	ttacgctaaa	taatcaacag	gtctgggttg	atagtcgcgc	tttgagcaca	5460
acaatcatg	aagccatgaa	tgataaatatg	tatgtaaata	gcagccaacg	gacagatggc	5520
tgttggttaa	acgcacctta	tacgatgagt	gggctggtga	gggctggtga	tacacgttca	5580
gctaattgggc	gctatgtcca	tatttcaaaa	gcttattcaa	acgaagtcgg	caatacatat	5640
tacttgacga	atttgaatgg	tcaaagcaca	tggattgaca	agcgggcgtt	tactgtgacc	5700
ttcgatcagg	tggtggcatt	aaatgcaacg	attgtggcac	gccaacgacc	agatgggatg	5760


```
tttaagacag caccatatgg tgaagcgggg gcgcagtttg tcgattatgt gacaaactat 5820
aaccagcaaa ccgtgccagt aacaaagcaa cattcagatg ctccaggggaa tcaatggtac 5880
ttagcgacag tgaatgggac acaatactgg attgatcaac ggtcattttc accagtagta 5940
acgaaggtgg ttgattatca agctaagatt gtgccacgga caacacgtga tgggtgtgtt 6000
agtggcgcac cctatgggga agtgaatgct aagctagtta acatggcaac tgcgtatcaa 6060
aatcaagttg tccatgcgac aggggaatat acgaatgctt cagggatcac atggagtcag 6120
ttcgcgttaa gcgggcaaga agacaagcta tggattgata agcgtgcttt gcaagcttaa 6180
gggaaggatt cgacaaaggt cgac                                     6204
```

<210> 21

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 21

Asp Thr Asn Ser Asn Val Ala
1 5

<210> 22

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 22

Met Lys Gln Gln Glu
1 5

<210> 23

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 23

Lys Lys Val Pro Val
1 5

<210> 24

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 24

Lys Asp Asp Glu Asn
1 5

<210> 25

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 25

Ile Asp Gly Asn Leu
1 5

<210> 26

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 26

Tyr Val Ala Asp Ser
1 5

<210> 27

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 27

His Leu Arg Lys Asn
1 5

<210> 28

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 28

Asn Glu Asn Thr Pro
1 5

<210> 29
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 29
Asn Val Asp Gly Tyr
1 5

<210> 30
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 30
Asn Pro Asp Leu Lys
1 5

<210> 31
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 31
Ser Asn Asp Ser Gly
1 5

<210> 32
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 32
Asn Thr Phe Val Lys
1 5

<210> 33
<211> 5

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 33
Ile Ser Gly Tyr Leu
1 5

<210> 34
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 34
Ser Asn Ala Ala Leu
1 5

<210> 35
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 35
Arg Gln Tyr Thr Asp
1 5

<210> 36
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 36
Gln Leu Tyr Arg Ala
1 5

<210> 37
<211> 5
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 37

Asp Asp Lys Ala Pro
1 5

<210> 38

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 38

Thr Arg Gln Tyr Thr
1 5

<210> 39

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 39

Ile Thr Phe Ala Gly
1 5

<210> 40

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 40

Asn Gln Tyr Lys Gly
1 5

<210> 41

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial sequence

<400> 41

Leu Phe Leu Asn Ala
1 5

<210> 42
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 42
Gln Val Ser Asp Thr
1 5

<210> 43
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 43
Leu Ile Thr Leu Asn
1 5

<210> 44
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 44
Gly Arg Tyr Val His
1 5

<210> 45
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 45
Thr Ala Pro Tyr Gly
1 5

<210> 46
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 46
Val Val Asp Tyr Gln
1 5

<210> 47
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 47
Leu Ser Gly Gln Glu
1 5

<210> 48
<211> 26
<212> DNA
<213> Artificial Sequence

<400> 48
tctagactgc aaaatggcaa ctacta 26

<210> 49
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: artificial
sequence

<400> 49
gtcgacgggtt tcatttggag tagtta 26

<210> 50
<211> 276
<212> DNA
<213> Solanum tuberosum

<400> 50
atggcaacta ctaaattcttt ttttaatttta ttttttatga tattagcaac tactagttca 60
acatgtgcta agttggaaga aatgggttact gttctaagta ttgatggagg tggaattaag 120
ggaatcattc cagctatcat tctcgaattt cttgaaggac aacttcagga agtggacaat 180
aataaagatg caagacttgc agattacttt gatgtaattg gaggaacaag tacaggaggt 240

ttattgactg ctatgataac tactccaaat gaaacc

276

<210> 51

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 51

gcgtactcta gacgtactcc gccatgacca c

31

<210> 52

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 52

gcgtacgtcg acggccctga tgggtcccat

30

<210> 53

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 53

ggccgggtcg acgatacaaa ttcgaatgtc gctg

34

<210> 54

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: artificial
sequence

<400> 54

ggccgggtcg aggttaccct cctttgtcga atc

33